

Supplementary file 1.

Crude 2013 point prevalence and cumulative incidence of gout (95% CI)

Variable	Number of persons in the 2013 population	Point prevalence (%)	95% CI	Number of persons at risk	Cumulative incidence per 10 000	95% CI
All	1 013 076	1.69	[1.66,1.71]	996 049	24	[23,25]
Women	515 154	0.96	[0.93,0.98]	509 432	15	[14,16]
Men	497 922	2.44	[2.40,2.49]	486 617	33	[32,35]
Age (years)						
18-34	288 270	0.08	[0.07,0.09]	286 780	2	[2,3]
35-44	167 363	0.43	[0.39,0.46]	167 101	7	[6,8]
45-54	166 197	1.04	[1.00,1.09]	162 838	14	[12,16]
55-64	147 060	1.90	[1.83,1.97]	146 969	26	[24,29]
65-74	135 727	3.53	[3.43,3.63]	129 274	55	[51,59]
75-84	74 122	5.87	[5.70,6.04]	70 425	82	[75,88]
85+	34 337	7.15	[6.88,7.43]	32 663	94	[83,105]
Education (years)						
0-9	227 552	2.85	[2.78,2.91]	231 528	38	[36,41]
10-12	400 004	1.73	[1.69,1.77]	403 678	24	[23,26]
13-14	121 781	1.21	[1.14,1.27]	123 782	16	[14,18]
15+	178 895	1.04	[1.00,1.09]	181 202	13	[12,15]
Income						
Low	232 553	0.92	[0.88,0.96]	226 356	13	[12,15]
Middle	484 488	2.23	[2.19,2.27]	490 407	30	[29,32]
High	247 704	1.64	[1.59,1.69]	248 016	23	[21,24]
Occupation						
White collar, high skilled	289 996	1.69	[1.64,1.74]	291 086	23	[21,25]
White collar, low skilled	221 339	1.21	[1.17,1.26]	223 635	18	[16,20]
Blue collar, high skilled	85 521	2.95	[2.84,3.07]	85 473	38	[34,43]
Blue collar, low skilled	123 379	2.16	[2.07,2.24]	124 018	28	[25,31]