



A. SAE incidence rates over time by indication in RCT+OLE patients




Indication	0-3	3-6	6-12	12-18	18-24	24-30	30-36	36-42	42-48	48-60	60-72	72-84	84-96
RA	321/6,927	253/6,581	344/5,743	212/4,003	184/2,782	123/2,410	105/1,817	104/1,702	98/1,511	144/1,296	49/914	31/249	2/64
axSpA	9/315	10/311	11/296	10/274	12/253	10/243	4/234	11/222	6/210	2/140	0/0	0/0	0/0
PsA	15/393	14/384	19/370	22/345	14/325	14/314	13/305	10/297	11/280	3/224	0/0	0/0	0/0
PSO	29/1,112	23/1,083	38/1,003	32/802	14/570	1/84	0/1	0/0	0/0	0/0	0/0	0/0	0/0
CD	283/2,570	167/2,388	178/1,741	120/1,188	91/913	64/746	43/624	27/485	19/364	47/282	35/227	19/184	9/155

B. TB disease incidence rates pre-2007 and from 2007 onwards over time in RA and CD RCT+OLE patients



	0-3	3-6	6-12	12-18	18-24	24-30	30-36	36-42	42-48	48-60	60-72	72-84	84-96
RA pre-2007	5/2,367	6/2,260	8/2,121	3/1,840	8/1,647	5/1,461	2/1,385	2/1,329	2/1,273	1/1,216	0/914	0/249	0/64
CD pre-2007	1/2,118	1/1,953	1/1,385	3/919	0/707	0/571	1/485	0/395	0/321	0/276	0/227	0/184	0/155
RA 2007 onwards	0/4,560	3/4,321	4/3,622	2/2,163	0/1,135	0/949	0/432	0/373	0/238	0/80	0/0	0/0	0/0
CD 2007 onwards	0/452	0/435	0/356	1/269	0/206	0/175	0/139	0/90	0/43	0/6	0/0	0/0	0/0

C. Malignancy incidence rates over time by indication in RCT+OLE patients



Indication	0-3	3-6	6-12	12-18	18-24	24-30	30-36	36-42	42-48	48-60	60-72	72-84	84-96
RA	7/6,927	14/6,581	26/5,743	19/4,003	16/2,782	9/2,410	10/1,817	8/1,702	7/1,511	11/1,296	6/914	1/249	0/64
axSpA	0/315	0/311	0/296	0/274	0/253	0/243	1/234	2/222	1/210	1/140	0/0	0/0	0/0
PsA	0/393	0/384	2/370	0/345	1/325	2/314	0/305	0/297	1/280	0/224	0/0	0/0	0/0
PSO	1/1,112	0/1,083	4/1,003	4/802	1/570	0/84	0/1	0/0	0/0	0/0	0/0	0/0	0/0
CD	4/2,570	1/2,388	5/1,741	2/1,188	6/913	2/746	2/624	2/485	1/364	5/282	2/227	2/184	1/155