Original research

Introduction and switching of biologic agents are associated with antidepressant and anxiolytic medication use: data on 42 815 real-world patients with inflammatory rheumatic disease

Vasiliki-Kalliopi Bournia ¹, Maria G Tektonidou ¹, Dimitrios Vassilopoulos,¹ Katerina Laskari,¹ Stylianos Panopoulos,¹ Kalliopi Fragiadaki,¹ Konstantinos Mathioudakis,² Anastasios Tsolakidis,² Panagiota Mitrou,³ Petros P Sfikakis¹

To cite: Bournia V.K, Tektonidou MG, Vassilopoulos D, *et al.* Introduction and switching of biologic agents are associated with antidepressant and anxiolytic medication use: data on 42 815 real-world patients with inflammatory rheumatic disease. *RMD Open* 2020;**6**: e001303. doi:10.1136/rmdopen-2020-001303

Received 4 May 2020
Revised 29 July 2020
Accepted 5 September 2020

© Author(s) (or their employer(s)) 2020. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to
Vasiliki-Kalliopi Bournia;
lily_bournia@hotmail.com and
Petros P Sfikakis;
psfikakis@med.uoa.gr

ABSTRACT

Objectives Depression and anxiety are linked bi-directionally with inflammatory rheumatic diseases (IRDs) activity, which in turn, depends on subjective patient reported outcomes that can be distorted by comorbid mood disorders. We tested the hypothesis that introduction and/or switching of biologic agents for IRDs are associated with treatment for depression and/or anxiety, by analysing real-world data.

Methods Using a country-wide electronic prescription database (10 012 604 registered, 99% population coverage), we captured almost all patients with rheumatoid arthritis (n=12 002), psoriatic arthritis (n=5465) and ankylosing spondylitis (n=6423) who received biologic disease modifying anti-rheumatic drugs (bDMARDs) during a 2-year period (8/2016–7/2018). Concomitant antidepressant/anxiolytic medication use was documented in patients who started or switched bDMARDs and compared with those who remained on conventional synthetic (cs)DMARDs or the same bDMARD, respectively, by multivariate regression analysis.

Results Two-year data analysis on 42 815 patients revealed that bDMARD introduction was associated with both antidepressant [OR: 1.248, 95% CI 1.153 to 1.350, p<0.0001] and anxiolytic medication use [OR: 1.178, 95% CI 1.099 to 1.263, p<0.0001]. Moreover, bDMARD switching was also associated with antidepressant [OR: 1.502, 95% CI 1.370 to 1.646, p<0.0001] and anxiolytic medication use [OR: 1.161, 95% CI 1.067 to 1.264, p=0.001]. Notably, all these associations were independent of age, gender, underlying disease diagnosis and concomitant glucocorticoid or csDMARD medication use.

Conclusion In real-world settings, both introduction and switching of bDMARDs in patients with IRDs were associated with the presence of mood disorders. Although a causal relationship is uncertain, the impact of depression and anxiety should always be considered by physicians facing the decision to introduce or switch bDMARDs in patients with active IRDs.

Key messages

What is already known about this subject?

► Depression and anxiety are common in patients with inflammatory rheumatic diseases (IRDs) and may negatively affect patient outcomes.

What does this study add?

► Analysis of data from 42 815 patients with rheumatoid arthritis, psoriatic arthritis and ankylosing spondylitis who received bDMARDs and/or csDMARDs between 8/2016 and 7/2018 revealed that both initiation and switching of bDMARDs were associated with the use of antidepressant and anxiolytic medications, independently of age, gender, underlying disease and concomitant glucocorticoid or csDMARD use.

How might this impact on clinical practice?

► Mood disorders that can either directly affect IRD activity and/or influence subjective patient-reported outcomes should be taken into account by rheumatologists who face the decision to initiate or switch bDMARDs.

INTRODUCTION

Depression and anxiety are common in patients with inflammatory rheumatic diseases (IRDs) influencing disease activity, quality of life and patient outcomes.¹ The relationship between mood disorders and inflammation seems to be bi-directional, as chronic pain and inflammation are considered to be important mediators of depression, while at the same time depression affects perception of pain and reduces response to treatment, possibly by

minimising patient adherence to medication.² In addition, inflammatory disease activity assessment depends largely on subjective patient reported outcomes that can be severely distorted by comorbid mood disorders, thus affecting the physicians' decisions.

Biologic disease modifying anti-rheumatic drugs (bDMARDs) were introduced for the treatment of patients with rheumatoid arthritis (RA) in 1998 and have ever since revolutionised the treatment of patients with chronic non-infectious inflammatory diseases who do not respond adequately to conventional treatment.^{3 4} Although there is an extensive literature on the interplay between inflammation and depression or anxiety, to the best of our knowledge, a possible association between the decision to initiate and/or switch bDMARDs for inflammatory arthritis and spondylitis and the presence of mood disorders has not been examined.

Herein, we had the opportunity to analyse big data on the use of bDMARDs for IRDs, derived from a country-wide electronic prescription database. Because the cost of bDMARDs is fully reimbursed by the national healthcare system, even for persons without insurance, all patients receiving these drugs among a country population of approximately 10 400 000 people are included in the database. An almost 100% population coverage and completeness of the data allowed us to test the hypothesis that introduction and/or switching of bDMARDs is associated with antidepressant and/or anxiolytic medication use in real-world patients with IRDs.

PATIENTS AND METHODS

In this retrospective population-based study, the e-Government Centre for Social Security Services prescription database (IDIKA) was searched using prespecified ICD-10 codes to capture every patient with RA (ICD-10 codes: M05, M05.0, M05.1, M05.2, M05.3, M05.8, M05.9, M06, M06.2, M06.3, M06.4, M06.8, M06.9), psoriatic arthritis (PsA, ICD-10 codes: M07, M07.0, M07.1, M07.3, L40.5) and ankylosing spondylitis (AS, ICD-10 codes: M45, M46, M46.0, M46.1, M46.8, M46.9, M07.2) in Greece who had filled at least one prescription for bDMARDs [Adalimumab (ATC5: L04AB04), Certolizumab-pegol (L04AB05), Etanercept (L04AB01), Golimumab (L04AB06), Infliximab (L04AB02), Abatacept (L04AA24), Anakinra (L04AC03), Rituximab (L01XC02), Secukinumab (L04AC10), Tocilizumab (L04AC07), Ustekinumab (L04AC05)] between August 1, 2016 and July 31, 2018. All patients who had filled at least two prescriptions for conventional synthetic (cs)DMARDs [Methotrexate (L01BA01 and L04AX03 for subcutaneous and oral formulations, respectively), Leflunomide (L04AA13), Hydroxychloroquine (P01BA02), Sulfasalazine (A07EC01), Ciclosporin (L04AD01), Azathioprine (L04AX01), D-penicillamine (M01CC01)] during the same time period, were also included.

Patients with 'other arthritis' (M13, M13.0, M13.1, M13.8, M13.9), enthesopathies (M76.9, M77, M77.8),

iridocyclitides (H20, H20.0, H20.1, H20.8, H20.9), arthropathy related to either Crohn's disease (M07.4) or ulcerative colitis (M07.5) and 'other enteropathic arthritis' (M07.6) were excluded from the study. Also, for those patients who had two or more of the above-mentioned ICD-10 diagnoses during the time period under study, only the more recent was taken into consideration for classification purposes.

The following data were recorded for all patients during the 2-year period: age at baseline, gender, use of bDMARDs, csDMARDs, glucocorticoids [betamethasone (H02AB01), methylprednisolone (H02AB04), prednisolone (H02AB06)], antidepressants [tricyclic antidepressants (N06AA), selective serotonin reuptake inhibitors (N06AB), non-selective monoamine oxidase inhibitors (N06AF), monoamine oxidase-a inhibitors (N06AG), other antidepressants (N06AX)] and anxiolytic medications [benzodiazepine derivatives (N05BA), Diphenylmethane derivatives (N05BB), Azaspirodecanedione derivatives (N05BE), Carbamates (N05BC), Dibenzobicyclo-octadiene derivatives (N05BD), Other anxiolytics (N05BX)].

Introduction of bDMARDs between 8/2016 and 7/2018 was documented for patients who had not filled a prior prescription for any bDMARD ever since records were being kept in the IDIKA database (January 2012). A switch was recorded every time a patient filled a prescription for a bDMARD with a different Anatomical Therapeutic Chemical Classification (ATC5) code from the directly preceding one, during the study time period. Discontinuation of bDMARDs was noted for patients who did not fill any bDMARD prescription after July 31, 2018.

Permission for use of anonymised data deposited in IDIKA was obtained from the Greek Ministry of Health following the approval of our formal request, according to the European legislation for General Data Protection Regulation (April 27, 2016) and the national laws (4600/2019, 4624/2019, 3892/10, 3418/2005). Due to regulations intended to protect trade secret, information on specific bDMARD prescription patterns, other than anti-tumour necrosis factor (TNF)/non-anti-TNF agents, cannot be shown.

Use of any antidepressant and/or anxiolytic drugs was compared between patients who initiated a bDMARD and those who remained on csDMARDs, as well as between patients who switched bDMARD and those who remained on the same bDMARD, using multivariate logistic regression analysis. The level of statistical significance was set at $p < 0.05$. The SPSS statistical software package (IBM Corp. Released 2017. IBM SPSS Statistics for Windows, Version 25.0.; IBM Corp., Armonk, New York) was used for all analyses.

RESULTS

A total of 42 815 patients with IRDs among 10 012 604 persons registered in the database (0.43%, 99.5% population coverage), of whom 18 925 were only on csDMARDs, were

captured (table 1). A total of 23 890 patients with IRDs in Greece initiated or continued bDMARDs during the 2-year period. Methotrexate and anti-TNF agents were the most commonly used csDMARDs and bDMARDs, respectively, across all disease and age groups in our population-based cohort. Distribution of patients treated with bDMARDs between RA (n=12 002), PsA (n=5465) and AS (n=6423) and their demographics (table 1) were comparable to our previously published country-wide prescription data for the period between 1/6/2014 and 31/5/2015.⁵

The number of patients with IRD who initiated, switched or discontinued a bDMARD between 1/8/2016 and 31/7/2018, stratified by diagnosis (RA, PsA, AS), age and gender, are shown in table 2. The percentage of patients that switched from one bDMARD to another was higher among PsA (18%) vs RA (13%) and AS (13.5%) patients (χ^2 test, $p<0.0001$), with women having more switches than men in all disease subgroups (RA group: 13.6% in females vs 10.2% in males, PsA group: 21.2% in females vs 14.3% in males, AS group: 50.5% in females vs 49.5% in males, χ^2 test, $p<0.0001$ for all comparisons).

The mean age of patients switching from one bDMARD to another was lower compared with non-switchers in the PsA group (54.96 $\pm$ 12.84 vs 56.20 $\pm$ 13.12 years, Student's *t*-test, $p=0.007$), marginally so in the RA group (63.06 $\pm$ 13.44 vs 63.80 $\pm$ 14.02 years, Student's *t*-test, $p=0.051$) and did not differ significantly in the AS group (50.79 $\pm$ 12.88 vs 51.24 $\pm$ 13.07 years, Student's *t*-test, $p=0.351$).

Table 3 shows the number of switches from one anti-TNF agent to another, or to a non-TNF bDMARD (abatacept, anakinra, rituximab, tocilizumab, secukinumab, ustekinumab) and vice-versa, in patients who filled bDMARD prescriptions for IRDs during the study period. Higher numbers of switches per patient were noted in RA (1.47) vs PsA (1.12) and AS (1.09).

Antidepressant and anxiolytic medication usage were documented in 24% and 43% of patients with RA, 19% and 36% of patients with PsA and in 16% and 30% of patients with AS, respectively. As shown in figures 1 and 2, anti-depressant and anxiolytic medication were more common among older patients in our cohort. Accordingly, after adjustment for age, gender, underlying disease and concomitant treatment with glucocorticoids, bDMARDs and csDMARDs, patients with AS had a higher probability [OR: 1.130, 95% CI 1.031 to 1.238, $p=0.009$], while patients with RA had a lower probability [OR: 0.880 95% CI 0.821 to 0.943, $p<0.0001$] to receive antidepressants compared with patients with PsA. Likewise, after adjustment for the same confounders, AS did not differ significantly from patients with PsA, whereas RA patients had a lower probability to use anxiolytics compared with patients with PsA [OR: 0.817, 95% CI 0.770 to 0.866, $p<0.0001$].

As also shown in the figures, antidepressant (figure 1) and anxiolytic (figure 2) usage were more frequent among all age and disease subgroups of patients who

initiated bDMARDs compared with those who remained on csDMARDs, and in any age, or disease subgroup of patients who switched to bDMARDs, compared with those who remained on the same therapy (with some exceptions in the subgroup of patients aged <45 years, for both antidepressants and anxiolytics). Multivariate analysis adjusted for age, gender, underlying disease and concomitant treatment with glucocorticoids and csDMARDs revealed a positive association between bDMARD introduction and usage of antidepressants [OR: 1.248, 95% CI 1.153 to 1.350, $p<0.0001$] or anxiolytics [OR: 1.178, 95% CI 1.099 to 1.263, $p<0.0001$]. Likewise, after correction for age, gender, underlying disease and concomitant treatment with glucocorticoids, csDMARDs and anti-TNF usage, multivariate analysis revealed a positive, independent association between bDMARD switching and antidepressant [OR: 1.502, 95% CI 1.370 to 1.646, $p<0.0001$] or anxiolytic medication usage [OR: 1.161, 95% CI 1.067 to 1.264, $p=0.001$].

DISCUSSION

To the best of our knowledge, this is the first study to show a positive association between the use of medications for depression or anxiety and both introduction and switching of bDMARDs, in a large, real-world population of patients with IRDs. Notably, these associations were independent of age, gender, underlying disease and concomitant glucocorticoid or csDMARD use.

Given the complex and bi-directional relationship of mood disorders with IRDs, the observed association may have at least three explanations, which are not mutually exclusive.

First, it has been previously shown that increased pain and physical disability, subsequent loss of social roles and employment, as well as medication side effects at higher disease activity, may predispose to depression.⁶ More severe inflammation corresponds to higher levels of circulating pro-inflammatory cytokines, which have been implicated in the pathogenesis of depression through a direct effect on brain structures known to contribute to the development of mood disorders.^{2,6} In fact, a recent large-scale study from Sweden indicated that in RA, PsA and AS patients, use of antidepressants and benzodiazepine-related agents decreased directly or soon after introduction of bDMARDs or non-biologics, possibly due to a collateral attenuation of psychiatric symptoms by the anti-inflammatory medications.⁷ The link between depression and inflammation is also supported by the fact that patients with major depressive disorder harbour increased levels of pro-inflammatory cytokines, such as TNF, interleukin (IL)-1 β and IL-6, increased cytokine receptors, acute phase reactants, such as C-reactive protein (CRP) and soluble adhesion molecules, both in the peripheral blood and in the cerebrospinal fluid. Interestingly, increased peripheral inflammatory markers were more common among anti-depressant non-responders compared with responders.⁸ Furthermore, it is well

Table 1 Country-wide data stratified by diagnosis and age on demographics and medication use, including switches, of real-world patients with inflammatory rheumatic disease who filled prescription for DMARDs between 8/2016 and 7/2018

	Rheumatoid arthritis (n=27 462)			Psoriatic arthritis (n=8469)			Ankylosing spondylitis (n=6883)		
	<45 years (n=1933)	45–65 years (n=10 317)	>65 years (n=15 212)	<45 years (n=1429)	45–65 years (n=4499)	>65 years (n=2541)	<45 years (n=2145)	45–65 years (n=3738)	>65 years (n=1000)
Female gender (%)	1576 (82)	8513 (83)	11 975 (79)	702 (49)	2422 (54)	1404 (55)	831 (39)	1500 (40)	376 (38)
csDMARDs									
Methotrexate (%)	1076 (56)	6172 (60)	8321 (55)	688 (48)	2474 (55)	1436 (57)	315 (15)	804 (22)	265 (27)
Leflunomide (%)	329 (17)	3130 (30)	5110 (34)	129 (9)	749 (17)	522 (21)	39 (2)	131 (4)	35 (4)
Hydroxychloroquine (%)	575 (30)	2846 (28)	3813 (25)	31 (2)	91 (2)	57 (2)	26 (1)	74 (2)	15 (2)
Other csDMARDs (%)	205 (11)	703 (7)	649 (4)	382 (27)	929 (21)	366 (14)	166 (8)	292 (8)	74 (7)
bDMARDs									
Anti-TNF (%)	934 (34)	3697 (36)	4230 (28)	890 (62)	2394 (53)	1095 (43)	1989 (93)	3273 (88)	872 (87)
Non-anti-TNF (%)	336 (17)	1554 (15)	2095 (14)	338 (24)	972 (22)	409 (16)	180 (8)	410 (11)	97 (10)
Glucocorticoids (%)	1073 (56)	6262 (61)	10 136 (67)	472 (33)	1544 (34)	1008 (40)	376 (18)	755 (20)	271 (27)
Anxiolytics (%)	323 (17)	3825 (37)	7788 (51)	226 (16)	1525 (34)	1281 (50)	309 (14)	1240 (33)	475 (48)
Antidepressants (%)	241 (12)	2394 (23)	3988 (26)	143 (10)	834 (19)	591 (23)	235 (11)	673 (18)	208 (21)

bDMARD, biologic DMARD; csDMARD, conventional synthetic DMARD; DMARD, disease modifying anti-rheumatic drugs; TNF, tumour necrosis factor.

Table 2 Country-wide data stratified by diagnosis, age and gender on patients who initiated, switched or discontinued (including deaths) any bDMARD, among all patients who filled at least one bDMARD prescription for inflammatory rheumatic disease between 8/2016 and 7/2018

	Rheumatoid arthritis (n=12 002)			Psoriatic arthritis (n=5465)			Ankylosing spondylitis (n=6423)		
	<45 years (n=1185)	45–65 years (n=4900)	>65 years (n=5917)	<45 years (n=1106)	45–65 years (n=2994)	>65 years (n=1365)	<45 years (n=2060)	45–65 years (n=3457)	>65 years (n=906)
Initiation									
Male (%)	83 (7)	281 (6)	310 (5)	193 (17)	370 (12)	134 (10)	406 (20)	471 (14)	119 (13)
Female (%)	347 (29)	1240 (25)	1246 (21)	195 (18)	514 (17)	204 (15)	348 (17)	451 (13)	84 (9)
Switching									
Male (%)	25 (2)	101 (2)	109 (2)	95 (9)	210 (7)	76 (6)	130 (6)	232 (13)	67 (7)
Female (%)	136 (11)	569 (12)	613 (10)	108 (10)	356 (12)	132 (10)	165 (8)	215 (6)	57 (6)
Discontinuation									
Male (%)	36 (3)	141 (3)	280 (5)	43 (4)	110 (4)	105 (8)	136 (7)	219 (6)	82 (9)
Female (%)	142 (12)	590 (12)	1013 (17)	58 (5)	188 (6)	116 (8)	140 (7)	200 (6)	58 (6)

bDMARD, biologic disease modifying anti-rheumatic drugs.

Table 3 Country-wide data stratified by diagnosis on bDMARD switches from an anti-TNF to a non-anti-TNF or to another anti-TNF and vice versa, in patients who filled bDMARD prescriptions for inflammatory rheumatic disease between 8/2016–7/2018

bDMARD switches	Rheumatoid arthritis (n=2290)	Psoriatic arthritis (n=1097)	Ankylosing spondylitis (n=948)
From anti-TNF to non-anti-TNF (%)	719 (31)	448 (41)	315 (33)
From one anti-TNF to another anti-TNF (%)	738 (32)	351 (32)	514 (54)
From a non-anti-TNF to an anti-TNF (%)	486 (21)	167 (15)	118 (12)
From non-anti-TNF to another non-anti-TNF (%)	347 (15)	131 (12)	1 (0.1)

Non-anti-TNF bDMARDs include: Abatacept, Anakinra, Rituximab, Tocilizumab, Secukinumab, Ustekinumab.

bDMARD, biologic disease modifying anti-rheumatic drugs; TNF, tumour necrosis factor.

known that patients who receive interferon- α therapeutically have a 15–40% risk to develop depression as a side effect, a rate significantly higher compared with untreated patients.⁹ It has been postulated that interferon- α , a potent cytokine inducer, triggers mood disorders by heightening right amygdala emotional reactivity. Conversely, treatment with anti-TNF agents has been shown to induce remission of underlying depression in the context of systemic autoimmune diseases, possibly by decreasing right amygdala reactivity.¹⁰ Other bDMARDs have also shown some effect in improving mental health in patients with systemic autoimmune diseases, with anti-IL6 therapy being the most effective.¹¹ It is

therefore possible that either the increased activity of the underlying IRD, and/or the increased levels of inflammation, could account for the more frequent use of antidepressant and anxiolytic medications among patients initiating or switching biologic agents.

A second possible explanation for the observed associations could be the fact that depression per se may induce or aggravate IRDs.¹² In a longitudinal cohort of psoriasis patients followed up for 25 years, major depressive disorder was an independent risk factor for the development of PsA, with a HR of 1.37.¹³ Furthermore, a significant association between major depressive disorder and subsequent development of RA has been shown in a cohort study from Taiwan [HR 1.65]¹⁴ and in a population-based cohort study from the UK [HR 1.38].¹⁵ In the later study, the risk for RA development was found to be attenuated by the use of antidepressants.¹⁵ In addition, a recent study showed that, according to patients' perception, psychological stress and mood disorders were the most frequent reasons underlying an RA flare.¹⁶ Patients with depression have 1.76 times higher odds for medication non-adherence compared with non-depressed patients,¹⁷ which could further account for the increased severity of co-morbid IRDs in this setting. Therefore, to the extent that depression can exacerbate IRDs, an association of antidepressant and anxiolytic medications with bDMARD initiation and switching can be anticipated.

A third explanation for the associations found between use of antidepressants or anxiolytics and the initiation or switching of bDMARDs in patients with IRD, could be that depression distorts subjective components of commonly used disease activity indices, thus misleading physicians to upscale treatment.^{8, 9} In a recent meta-analysis of patients with axial spondylarthritis reporting a 15% pooled prevalence of at least moderate depression, patients with depression displayed significantly worse disease activity (BASDAI, ASDAS) and functional impairment (BASFI, BASMI) indices compared with those without depression.

Figure 1 Percentages of antidepressant medication use among real-world patients with inflammatory rheumatic disease, stratified by age, diagnosis and by introduction (panel A) or switching (panel B) of biologic disease modifying anti-rheumatic drugs (bDMARDs). N denotes patient numbers per disease/age groups.

Figure 2 Percentages of anxiolytic medication use among real-world patients with inflammatory rheumatic disease, stratified by age, diagnosis and by introduction (panel A) or switching (panel B) of biologic disease modifying anti-rheumatic drugs (bDMARDs). N denotes patient numbers per disease/age groups.

Although a causal effect of higher disease activity on depression cannot be excluded, it is also possible that depression distorted perception of pain in patients with axial spondylarthritis. In fact, the meta-analysis showed that differences in subjective indices (BASDAI, BASFI) were larger than differences in objective measures (BASMI, ASDAS), possibly indicating that patients with depression tended to overrate pain and discomfort caused by their underlying rheumatic disease and therefore overestimate disease severity.¹⁰ Likewise, it has been shown that in patients with severe RA the patient visual analogue scale, one of the subjective components of DAS28, correlates with patients' beliefs regarding illness and treatment as well as with depression.¹¹ The influence of baseline depression and anxiety on the subjective components of DAS28 at 1-year follow-up is further supported by the results of a small study including 56 patients with RA, which showed that tender joint count and patient global assessment were the two DAS-28 parameters particularly affected by mood disorders.⁹ Furthermore, in a prospective observational study including 1326 RA and 728 PsA patients, baseline depression or anxiety was associated with increased subjective measures of disease activity, such as patient's and physician's global assessment and with tender joint count and joint pain in RA at follow-up, but not with objective measures such as swollen joint count and acute phase reactants.⁸

In addition to the associations discussed above, the analysis presented herein revealed that higher rates of antidepressant medication usage occur in AS, followed by PsA patients, while RA patients had the lowest rates, after correction for age, gender, use of bDMARDs, csDMARDs and glucocorticoids, the latter of which may be per SE associated with mood disorders.¹⁸ Likewise, patients with AS and PsA showed higher rates of anxiolytic usage compared with patients with RA. So far, meta-analyses and cohort studies have shown higher odds for depression among patients with RA, PsA and AS^{19–22}

compared with the general population. Depression in rheumatic diseases has been associated with impaired health-related quality of life,²³ worse response to treatment and lower improvement in disease activity over time.^{8 12 24} Only few small cohort studies have directly compared the OR of depression between different IRDs, reporting an equal probability between patients with AS and RA²⁵ and a higher probability in patients with PsA compared with RA.²⁶ The higher burden of depression and anxiety among patients with AS and PsA compared with patients with RA found in our study conforms to some extent to the results of a recent study showing that disease burden as assessed by patient global assessment, pain and fatigue in PsA and AS was comparable to or greater than that in patients with RA.²⁷

In our previous analysis of data from the country-wide prescription database between 06/2014 and 05/2015, more than 99% of patients with AS and PsA (combined group) received an anti-TNF agent, whereas only 0.78% used a non-anti-TNF bDMARD. In the current study, the rate of anti-TNF usage among bDMARD treated patients with AS and PsA has fallen to 88% and the use of non-anti-TNF bDMARDs has risen to 20%. The fact that other bDMARDs approved for the indications of AS and PsA, such as ustekinumab or secukinumab, only became available around 2016, could have precipitated switching from an anti-TNF agent to a non-TNF bDMARD in these patient groups during the time period we are studying. However, as shown in table 3, switching from one anti-TNF agent to another, still represents about 42% of the total switches recorded.

In the 2-year period between 1/8/2016 and 31/7/2018 several biosimilars of anti-TNF agents have been marketed and many physicians switched their patients to these new drugs in an effort to cut down on medical costs. To avoid recording switching to biosimilars as a switch to another bDMARD, we searched the database using ATC5 codes for all medications of interest, given that the ATC classification system relies

on substance and cannot discriminate between a bDMARD and its biosimilar.

Our study has several limitations. First, we were not able to capture all patients treated with csDMARDs in our country. The reason is that a significant number of patients with IRDs had no relevant prescription records kept in the electronic database system and as a result, have not been included in our cohort, because they prefer to pay the very low cost of methotrexate and hydroxychloroquine in order to avoid the time-consuming process of having them prescribed, which in addition costs more than the drug prices for those followed in private units. Second, we were unable to retrieve reliable individual-level data on IRD activity and patient functional impairment due to the structure of the e-Government Centre for Social Security Services prescription database. Although disability is a factor possibly predisposing to anxiety and depression, data on disease duration that could correlate with disability was also not available. Third, our findings did not rely on a psychiatric diagnosis of depression and anxiety, but the diagnosis of mood disorders was indirectly inferred from the use of antidepressant and/or anxiolytic medications. As a result, the lack of data on the possible concomitant presence of fibromyalgia for which antidepressants and anxiolytics are commonly prescribed may confound our results. On the other hand, not all patients with RA or PsA and concomitant depression use antidepressants,^{20–28} therefore our approach has clearly missed some of them. Although a comparison of our findings with antidepressant and anxiolytic prescription patterns in a non-IRD population would be revealing, it would require data for which we have not requested access. Finally, our study examines the use of antidepressant and anxiolytic medications and associates them with the introduction and switching of bDMARDs in patients with IRD over the entire time span of the two-year period under study. Consequently, an exact temporal association of bDMARD introduction or switching with the use of psychiatric medications cannot be supported, mainly due to the lack of an index event (introduction or switching) in the control group.

A universal healthcare system with single-payer medication coverage and no out-of-pocket cost for bDMARDs, provides a unique opportunity to test our initial hypothesis on a population level, avoiding any selection bias. A major strength of our study is the fact that the database from which our cohort is derived covers almost the entire country's population, therefore practically all patients with RA, PsA and AS treated with bDMARDs in Greece. The large sample size allows for robust estimates of association, whereas current demographic findings are consistent with previous observations in the same real-world population.⁵

To conclude, depression and anxiety that are bi-directionally related with IRD, may contribute to bDMARD initiation and switching in these patients, either by aggravating disease severity and/or by distorting

the perception of patient-reported outcome measures. Mood disorders should be always considered by practicing rheumatologists upon the decision to propose to an 'inadequately treated' patient the introduction or switching of a biologic agent. Whether the same holds true for patients with psoriasis and inflammatory bowel disease needs further investigation.

Author affiliations

¹Joint Rheumatology Program, Medical School, National and Kapodistrian University of Athens, Athens, Greece

²DIKA SA - e-Government Center for Social Security Services, Athens, Greece

³Hellenic Republic Ministry of Health, Athens, Greece

Contributors VKB contributed in designing the study, statistical analysis, curation of data and drafting of the manuscript, MGGT contributed in designing the study and critically revised the manuscript, DV contributed in study conceptualisation, study design and critical revision of the manuscript, KL contributed in analysis of data, SP and KF contributed in data curation and drafting the manuscript. KM, AT and PM contributed in data curation and analysis, PPS conceived the original idea, supervised the project and contributed in study design, data analysis, writing and reviewing the manuscript.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement All data relevant to the study are included in the article.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Vasiliki-Kalliopi Bournia <http://orcid.org/0000-0002-2162-1362>

Maria G Tektonidou <http://orcid.org/0000-0003-2238-0975>

REFERENCES

- 1 Dregan A, Matcham F, Harber-Aschan L, *et al*. Common mental disorders within chronic inflammatory disorders: a primary care database prospective investigation. *Ann Rheum Dis* 2019;78:688–95.
- 2 Nerurkar L, Siebert S, McInnes IB, *et al*. Rheumatoid arthritis and depression: an inflammatory perspective. *Lancet Psychiatry* 2019;6:164–73.
- 3 Sarzi-Puttini P, Ceribelli A, Marotto D, *et al*. Systemic rheumatic diseases: from biological agents to small molecules. *Autoimmun Rev* 2019;18:583–92.
- 4 Sfikakis PP. The first decade of biologic TNF antagonists in clinical practice: lessons learned, unresolved issues and future directions. *Curr Dir Autoimmun* 2010;11:180–210.
- 5 Sfikakis PP, Bournia VK, Sidiropoulos P, *et al*. Biologic treatment for rheumatic disease: real-world big data analysis from the Greek country-wide prescription database. *Clin Exp Rheumatol* 2017;35:579–85.
- 6 Vallerand IA, Patten SB, Barnabe C. Depression and the risk of rheumatoid arthritis. *Curr Opin Rheumatol* 2019;31:279–84.
- 7 Brenner P, Citarella A, Wingård L, *et al*. Use of antidepressants and benzodiazepine-related hypnotics before and after initiation of TNF- α inhibitors or non-biological systemic treatment in patients with rheumatoid arthritis, psoriatic arthritis or ankylosing spondylitis. *BMC Rheumatol* 2020;4:9.
- 8 Michelsen B, Kristianslund EK, Sexton J, *et al*. Do depression and anxiety reduce the likelihood of remission in rheumatoid arthritis and psoriatic arthritis? Data from the prospective multicentre NOR-DMARD study. *Ann Rheum Dis* 2017;76:1906–10.
- 9 Matcham F, Ali S, Irving K, *et al*. Are depression and anxiety associated with disease activity in rheumatoid arthritis? A prospective study. *BMC Musculoskelet Disord* 2016;17:155.

- 10 Zhao S, Thong D, Miller N, *et al.* The prevalence of depression in axial spondyloarthritis and its association with disease activity: a systematic review and meta-analysis. *Arthritis Res Ther* 2018;20:140.
- 11 Cordingley L, Prajapati R, Plant D, *et al.* Impact of psychological factors on subjective disease activity assessments in patients with severe rheumatoid arthritis. *Arthritis Care Res (Hoboken)* 2014;66:861–8.
- 12 Matcham F, Norton S, Scott DL, *et al.* Symptoms of depression and anxiety predict treatment response and long-term physical health outcomes in rheumatoid arthritis: secondary analysis of a randomized controlled trial. *Rheumatology (Oxford)* 2016;55:268–78.
- 13 Lewinson RT, Vallerand IA, Lowerison MW, *et al.* Depression is associated with an increased risk of psoriatic arthritis among patients with psoriasis: a population-based study. *J Invest Dermatol* 2017;137:828–35.
- 14 Lu M-C, Guo H-R, Lin M-C, *et al.* Bidirectional associations between rheumatoid arthritis and depression: a nationwide longitudinal study. *Sci Rep* 2016;6:20647.
- 15 Vallerand IA, Lewinson RT, Frolkis AD, *et al.* Depression as a risk factor for the development of rheumatoid arthritis: a population-based cohort study. *RMD Open* 2018;4:e000670.
- 16 Yilmaz V, Umay E, Gündoğdu İ, *et al.* Rheumatoid arthritis: are psychological factors effective in disease flare? *Eur J Rheumatol* 2017;4:127–32.
- 17 Grenard JL, Munjas BA, Adams JL, *et al.* Depression and medication adherence in the treatment of chronic diseases in the United States: a meta-analysis. *J Gen Intern Med* 2011;26:1175–82.
- 18 Brown ES. Effects of glucocorticoids on mood, memory, and the hippocampus. *Ann N Y Acad Sci* 2009;1179:41–55.
- 19 Wang S-L, Chang C-H, Hu L-Y, *et al.* Risk of developing depressive disorders following rheumatoid arthritis: a nationwide population-based study. *PLoS One* 2014;9:e107791.
- 20 Zusman EZ, Howren AM, Park JYE, *et al.* Epidemiology of depression and anxiety in patients with psoriatic arthritis: a systematic review and meta-analysis. *Semin Arthritis Rheum*.
- 21 Meesters JJJ, Bremder A, Bergman S, *et al.* The risk for depression in patients with ankylosing spondylitis: a population-based cohort study. *Arthritis Res Ther* 2014;16:418.
- 22 Park JY, Howren AM, Zusman EZ, *et al.* The incidence of depression and anxiety in patients with ankylosing spondylitis: a systematic review and meta-analysis. *BMC Rheumatol* 2020;4:12.
- 23 Haugeberg G, Michelsen B, Kavanaugh A. Impact of skin, musculoskeletal and psychosocial aspects on quality of life in psoriatic arthritis patients: a cross-sectional study of outpatient clinic patients in the biologic treatment era. *RMD Open* 2020;6:e001223.
- 24 Matcham F, Davies R, Hotopf M, *et al.* The relationship between depression and biologic treatment response in rheumatoid arthritis: an analysis of the British society for rheumatology biologics register. *Rheumatology* 2018;57:835–43.
- 25 Hyphantis T, Kotsis K, Tsifetaki N, *et al.* The relationship between depressive symptoms, illness perceptions and quality of life in ankylosing spondylitis in comparison to rheumatoid arthritis. *Clin Rheumatol* 2013;32:635–44.
- 26 Sinnathurai P, Buchbinder R, Hill C, *et al.* Comorbidity in psoriatic arthritis and rheumatoid arthritis. *Intern Med J* 2018;48:1360–8.
- 27 Mease PJ, Liu M, Rebello S, *et al.* Comparative disease burden in patients with rheumatoid arthritis, psoriatic arthritis, or axial spondyloarthritis: data from two corona registries. *Rheumatol Ther* 2019;6:529–42.
- 28 Englbrecht M, Alten R, Aringer M, *et al.* New insights into the prevalence of depressive symptoms and depression in rheumatoid arthritis – implications from the prospective multicenter VADERA II study. *PLoS One* 2019;14:e0217412.