

**Table S2. Signs and symptoms identified in the targeted literature review**

<b>Disease-related signs/symptoms</b>	<b>No. of article mentions (n=15)</b>	<b>Reported prevalence range</b>	<b>Symptom salience*</b>
Pain (skin, joint, muscle, spine)	14 (93%)	65%–95%	High
Fatigue	12 (80%)	57%–84%	High
Skin symptoms (itching, red, scaly, etc)	9 (60%)	25%-76%	High
Stiffness	7 (47%)	77%	High
Swollen / Inflamed joints	7 (47%)	--	High
Gastrointestinal distress	2 (13%)	12%-75%	High
Nail symptoms (pitting, lifting)	6 (40%)	21%	Intermediate
Dactylitis (“sausage fingers”)	5 (33%)	45%	Intermediate
Musculoskeletal symptoms	5 (33%)	--	Intermediate
Impaired grip strength	5 (33%)	--	Intermediate
Deformed joints	3 (20%)	--	Intermediate
Respiratory symptoms	2 (13%)	19%	Intermediate
Difficulty swallowing / biting	2 (13%)	--	Low
Impaired vision	2 (13%)	--	Low
Limited dexterity	2 (13%)	--	Low
Paresthesia	1 (2%)	--	Low

\*High corresponded to prevalence higher than 75% and/or concept mentioned in >50% of articles; intermediate corresponded to prevalence of 30-74% and/or mentioned in 30-50% of articles; low corresponded to prevalence below 30% and/or mentioned in <30% of articles.