Correction: High levels of immunosuppression are related to unfavourable outcomes in hospitalised patients with rheumatic diseases and COVID-19: first results of ReumaCoV brasil registry

Marques CDL, Kakehasi AM, Pinheiro MM, *et al.* High levels of immunosuppression are related to unfavourable outcomes in hospitalised patients with rheumatic diseases and COVID-19: first results of ReumaCoV Brasil registry. *RMD Open* 2021;7:e001461. doi: 10.1136/rmdopen-2020-001461

The article has been corrected since it was published online. The authors want to alert the readers on missing to include co-author Felipe Omura in the authors' list. He has been identified one of the authors who had a vital contribution towards the preparation of the paper.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See <http://creativecommons.org/licenses/by-nc/4.0/>.

© Author(s) (or their employer(s)) 2021. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

RMD Open 2021;7:e001461corr1. doi:10.1136/rmdopen-2020-001461corr1