SHORT REPORT

High p16^{INK4a}, a marker of cellular senescence, is associated with renal injury, impairment and outcome in lupus nephritisGaëlle Tilman,^{1,2} Caroline Bouzin,³ Selda Aydin,⁴ Farah Tamirou,² Christine Galant,⁴ Pierre G Coulie,^{5,6} Frédéric Houssiau,^{2,7} Bernard Lauwerys,^{2,7} Nisha Limaye ¹

To cite: Tilman G, Bouzin C, Aydin S, *et al.* High p16^{INK4a}, a marker of cellular senescence, is associated with renal injury, impairment and outcome in lupus nephritis. *RMD Open* 2021;7:e001844. doi:10.1136/rmdopen-2021-001844

► Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/rmdopen-2021-001844>).

Received 26 July 2021
Accepted 5 October 2021

ABSTRACT

Objectives Because a significant fraction of patients with lupus nephritis (LN) develops renal impairment, there is a need to better understand the mechanisms underlying disease progression. Here, we assessed for cellular senescence in the LN kidney, and its association with disease severity and outcome.

Methods We enumerated the number of cells positive for p16^{INK4a} protein, a marker of cellular senescence, by immunohistochemistry followed by digital quantification, on renal biopsies from 40 patients with active LN. We tested for an association of p16^{INK4a} with renal fibrosis, CD8⁺ T cell infiltration, systemic disease and renal function at baseline and at 5 years.

Results The presence of p16^{INK4a}-positive cells was significantly associated with lower estimated glomerular filtration rate at baseline and 5 years post-treatment, independently of patient demographics and systemic disease parameters. It was also associated with higher baseline renal fibrosis and CD8⁺ T cell infiltration. Interestingly, we observed marked spatial co-distribution of glomerular p16^{INK4a}-positive cells with CD8⁺ T cells.

Conclusion We demonstrate, for the first time, that LN biopsies characterised by renal impairment display increased p16^{INK4a}-positive cells, associated with higher fibrosis and CD8⁺ T cell infiltration. Cellular senescence may represent a kidney-intrinsic disease mechanism and potentially, a novel therapeutic target in LN.

INTRODUCTION

Lupus nephritis (LN) is a severe complication of systemic lupus erythematosus (SLE), initiated by deposition of immune complexes or autoantibodies in glomerular basal membrane, followed by recruitment of inflammatory cells.¹ Renal injury leads to irreversible fibrosis, resulting in loss of kidney function. LN is treated with high-dose corticosteroids and other immunosuppressive agents.² One-third of patients nevertheless show a decline in renal function, with 5%–10% developing end-stage renal disease

Key messages

What is already known about this subject?

- Cellular senescence has been observed in renal ageing, certain renal diseases and in a mouse model of lupus nephritis, and was associated with renal damage.

What does this study add?

- This is the first demonstration of the occurrence of cellular senescence, based on the presence of p16^{INK4a}-positive cells, in a large series of lupus nephritis kidney biopsies, taken at baseline.
- The presence of p16^{INK4a}-positive cells is associated with renal fibrosis, CD8⁺ T cell infiltration and functional impairment, but not with patient demographics or systemic disease parameters.

How might this impact on clinical practice?

- The striking codistribution of p16^{INK4a}-positive cells and CD8⁺ T cells suggests a functional, interactive role in the pathogenesis of lupus nephritis, which may be targeted by senolytic therapies.
- The association with poor renal function 5 years post-treatment suggests the cellular senescence marker p16^{INK4a} in baseline renal biopsies may warrant exploration as a prognostic marker for poor renal outcome.

within 10 years.³ Prognostic markers that would allow for timely treatment escalation or modification are hence eagerly sought, as are novel therapeutic targets.

Cellular senescence, triggered by stimuli such as telomere erosion, oxidative stress and chronic inflammation, ultimately leads to irreversible growth arrest through the accumulation of cyclin dependent kinase (CDK) inhibitors including p16^{INK4a} protein (*CDKN2A*): a major hallmark of cellular senescence.⁴ Senescent cells remain metabolically active and undergo a number of morphological and physiological changes including the

© Author(s) (or their employer(s)) 2021. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Dr Nisha Limaye;
nisha.limaye@uclouvain.be

upregulation of β -galactosidase activity⁵ and acquisition of a proinflammatory, profibrotic senescence-associated secretory phenotype (SASP).⁶ While essential in tissue repair and remodelling (e.g., during embryogenesis), cellular senescence can exert adverse effects in aging-related and chronic disease as well as cancer.⁴ p16^{INK4a} or β -galactosidase positive cells have been observed in renal ageing and certain kidney diseases, and were associated with histological lesions and renal impairment.⁷ The presence of β -galactosidase positive cells correlated with proteinuria in MRL/*lpr* lupus-prone mice⁸; however, cellular senescence is yet to be clearly demonstrated in LN. Here, we report the occurrence of p16^{INK4a}-positive cells in kidney biopsies from (n=40) patients with active LN, and its association with renal injury and functional impairment.

METHODS

Patients and kidney biopsies

Patients were recruited at the Department of Rheumatology, Cliniques universitaires Saint-Luc (Brussels, Belgium). All met the 1982 revised ACR classification criteria for SLE and had biopsy-proven LN. Formalin-fixed paraffin-embedded (FFPE) renal biopsies were residual corporal material collected for diagnostic purposes between January 1996 and November 2019. Estimated glomerular filtration rate (eGFR) values were calculated using the Chronic Kidney Disease Epidemiology Collaboration (CKD-EPI) formula. Patient consent was not required for the use of residual corporal material, in agreement with Belgian regulations on human studies.

Histology and immunohistochemistry

ISN/RPS 2003 classification of SLE renal biopsies, semi-quantitative sclerosis scores, and National Institutes of Health (NIH) activity index (AI) and chronicity index (CI) were retrieved from medical records. Immunostaining with anti-CD8 (C8/144B, Dako) and anti-p16^{INK4a} (E6H4, Roche Ventana CIntec Histology) and Picosirius Red (PSR) staining were performed on 5 μ m FFPE serial sections. Slides were digitalised on an SCN400 scanner (Leica Biosystems, Germany) or a Panoramic Confocal slide scanner (3DHitech, Hungary) at $\times 20$ magnification. Computer-assisted quantification of the entire surface of sections was performed using AuthorTM V.2017.2 (Visiopharm, Denmark). Results shown are the number of p16^{INK4a}- or CD8-positive cells per μ m² of tissue. Semi-quantitative PSR scores (scale: 2–6) are median (glomerular + interstitial fibrosis) scores from three (blinded) scorers. Further details are provided in online supplemental methods 1, online supplemental figure 1.

Statistical analyses

Statistical analyses were performed on GraphPad Prism V.9.1.0: Mann-Whitney or Wilcoxon matched-pair signed rank tests for two-group comparisons, Kruskal-Wallis for multigroup comparisons and Spearman's rank-order correlation coefficient.

RESULTS

p16^{INK4a}-positive cells in kidney biopsies from patients with active LN

We evaluated p16^{INK4a} protein by immunohistochemistry on renal biopsies taken at diagnosis from 40 patients with active LN, including incident nephritis (n=31) and relapse (n=9). Demographic, biological and clinical data are summarised in online supplemental table 1. p16^{INK4a}-positive cells were detected in all, but with considerable variability between samples: from virtually none, to occasional scattered cells, to strongly positive areas (figure 1A, online supplemental figure 2A–D). Quantification of p16^{INK4a} staining confirmed the heterogeneity of LN biopsies (range: 1.76×10^{-6} – 260×10^{-6} , median: 14.7×10^{-6} cells/ μ m²) (figure 1B,C). Stained cells included mesangial cells, endothelial cells or podocytes in glomeruli, parietal epithelial cells in Bowman's capsules, proximal or distal tubular cells, and interstitial cells (figure 1D). Although the density of p16^{INK4a}-positive cells (per μ m²) was significantly higher in glomeruli than in interstitia (figure 1C), these values (per sample) showed a significant positive correlation ($r=0.7591$, $p<0.0001$) (online supplemental figure 3). Importantly, p16^{INK4a} accumulation was not associated with patient age, gender or ethnicity (online supplemental figure 4A–C).

p16^{INK4a} is associated with renal impairment at baseline and 5 years post-treatment

p16^{INK4a} positivity was significantly associated with impaired renal function: biopsies from patients with eGFR <60 at the time of sampling had significantly higher p16^{INK4a}-positive cells/ μ m², and conversely, samples with high (>75th percentile) p16^{INK4a} were associated with significantly lower baseline eGFR (figure 2A,B). The association with eGFR was true for both glomerular and interstitial p16^{INK4a} (online supplemental figure 5A,B). Interestingly, whereas high glomerular p16^{INK4a} was also associated with significantly higher proteinuria (urinary protein/creatinine ratio), which mainly reflects glomerular injury, interstitial p16^{INK4a} was not (online supplemental figure 5C,D). In contrast to its association with poor renal function, p16^{INK4a} was not associated with parameters of systemic disease such as serum anti double-stranded DNA antibody or C3 levels, or with ISN/RPS classification (online supplemental figure 6A–D). Slightly (non-significantly) higher p16^{INK4a} was observed in biopsies from patients with longer duration between SLE and LN diagnosis, and in relapse compared with incident nephritis (online supplemental figure 7A,B). Importantly, analysis of only the incident LN cases upheld the association between p16^{INK4a} and eGFR (online supplemental figure 7C), suggesting it is not solely dependent on kidney disease duration. Finally, high p16^{INK4a} in baseline biopsies was also associated with lower eGFR at 5 years post-treatment (but not at 1 year), suggesting it may be predictive of poor long-term renal evolution (figure 2C,D).

Figure 1 p16^{INK4a}-positive cells in kidney biopsies from patients with active LN. (A) p16^{INK4a}, CD8 and PSR staining of FFPE serial sections from an LN kidney baseline biopsy. Overview (left panels), close-up (right) showing p16^{INK4a}-positive cells within a glomerulus, periglomerular CD8-positive cells and collagen deposits in and around Bowman's capsule of the same glomerulus. (B) Quantification of p16^{INK4a} (positive cells/μm²) showing degree of heterogeneity among samples. (C) Quantification of p16^{INK4a} (positive cells/μm²) in glomerular vs interstitial areas of the same samples. *p* value: Wilcoxon matched-pair signed rank test. (D) Representative image from a sample showing p16^{INK4a}-positive cells within glomeruli (#), in Bowman's capsules (filled star), proximal tubules (filled arrow), distal tubules (arrow) and dispersed in interstitium (arrowhead). FFPE, formalin-fixed paraffin-embedded; LN, lupus nephritis; PSR, picrosirius red.

p16^{INK4a} is associated with renal CD8⁺ T cell infiltration and fibrosis

It has been shown that CD8⁺ T lymphocytes are the predominant immune cell type infiltrating the LN kidney,⁹ and that their presence is associated with renal disease severity.^{9,10} We found that p16^{INK4a}-high biopsies displayed significantly higher CD8⁺ T cell infiltration (figure 3A). They were also significantly associated with increased fibrosis: collagen

Figure 2 Association between p16^{INK4a}-positive cells and renal impairment in LN. (A) Quantification of p16^{INK4a} (positive cells/μm²) in samples from patients with renal impairment (estimated glomerular filtration rate: eGFR <60 mL/min/1.73 m²) at baseline. (B–D) eGFR in patients with p16^{INK4a}-high (>75th percentile, 'Q3') vs p16^{INK4a} low-to-moderate (<Q3) baseline kidney biopsies. Decreased eGFR at baseline (significant) (B) 1-year (non-significant) (C) and 5 years post-treatment (significant) (D) in p16^{INK4a}-high vs p16^{INK4a} low-to-moderate group. Horizontal bars: medians. *p* values: Mann-Whitney U test.

deposition as reflected by semi-quantitative scores of PSR staining, and sclerosis and NIH CI scores provided by a nephropathologist (figure 3B–D). NIH AI scores, in contrast, were not significantly different between the p16^{INK4a} groups (data not shown). Intriguingly, in examining tissue sections, we noticed what seemed to be a spatial co-distribution between strong p16^{INK4a} staining (that tended to be glomerular) and clustered CD8-positive cells (most often periglomerular) in certain LN biopsies (figure 1A, online supplemental figure 2). We, therefore, quantified the number of CD8⁺ T cells located within a 30 μm radius around each glomerulus, the thickness of its Bowman's capsule, and the number of p16^{INK4a}-positive cells within it (online supplemental figure 1). This was done on all glomeruli visible on the three serial sections, across all samples. Glomeruli with high p16^{INK4a}-positive cell accumulation displayed significantly higher periglomerular CD8⁺ T cells (figure 3E), as well as increased thickness of Bowman's capsules, reflecting glomerular fibrosis (figure 3F).

Figure 3 Association between p16^{INK4a}-positive cells, CD8⁺ T cell infiltration and fibrosis in LN kidney biopsies. (A) Significantly higher CD8⁺ T cells (positive cells/μm²) in p16^{INK4a}-high (>75th percentile, “Q3”) vs p16^{INK4a} low-to-moderate (<Q3) biopsies. (B–D) Significantly higher fibrosis in p16^{INK4a}-high vs p16^{INK4a} low-to-moderate biopsies: collagen deposits (semi-quantitative PSR staining scored on a scale from 2 to 6) (B), sclerosis (glomerulosclerosis and interstitial fibrosis on a scale from 0 to 6) (C), and NIH chronicity index (on a scale from 0 to 12) (D). (E, F) Glomeruli with high p16^{INK4a} (>Q3) show significantly higher numbers of periglomerular CD8⁺ T cells (quantified in a 30 μm radius around the glomerulus) (E), and significantly thicker Bowman’s capsule (fibrous tissue surrounding the capsule, normalised to area of the glomerulus) (F). Horizontal bars: medians. *p* values: Mann-Whitney U test. PSR, picrosirius red.

DISCUSSION

This is the first demonstration, in a large series of patients, that p16^{INK4a}, a major hallmark of cellular senescence, is associated with disease severity in LN. IHC for p16^{INK4a} protein is widely used to assess for senescence *ex vivo*, and has been associated with severity of other renal diseases.⁷ Selective elimination of p16^{INK4a}-positive cells in mouse models of ageing and induced nephropathy has moreover been shown to relieve fibrotic lesions and improve renal function,^{11 12} suggesting a key role in renal pathogenesis. Why cellular senescence may occur in LN is unknown. The inflammatory, oxidative environment of the LN kidney may be a source of cell stress. Several proinflammatory factors have been implicated in senescence induction,^{13 14} including interferon-β¹³ that has been linked to the senescent phenotype of bone marrow mesenchymal stem cells from patients with SLE.¹⁵

While our study highlights a significant association between the abundance of p16^{INK4a}-positive cells and LN severity, the use of additional markers of cellular senescence, in a larger cohort, will be essential to confirm its relevance in LN. If and how cellular senescence contributes to disease progression (or whether it simply reflects

tissue injury) remains to be investigated. A detrimental effect may be exerted through the profibrotic, proinflammatory secretome (SASP) typical of senescent cells. In keeping with this hypothesis, we describe a tight spatial co-distribution between p16^{INK4a} positive cells, fibrosis and CD8⁺ T cell infiltration in LN kidneys. Another pathogenic mechanism may involve the accumulation of functionally incompetent cells, for instance, renal progenitor cells (RPC, a subset of parietal epithelial cells located in the Bowman’s capsule of glomeruli).¹⁶ While healthy RPCs regenerate glomerular and tubular structures thanks to their capacity to proliferate and differentiate into renal cell subsets,¹⁶ senescent RPCs could hamper tissue repair.¹⁷ Finally, while the SASP is particularly suited to engaging the immune system (including CD8⁺ T lymphocytes)¹⁸ for the clearance of senescent cells, the latter can upend the process by inhibiting cytolytic cells.¹⁸ It has been suggested that persistence of senescent cells (due to the overwhelming or inhibition of the immune response) tips the balance from a positive to a negative impact.¹⁶ Further *in vitro* experiments will be required in order to dissect the role of p16^{INK4a}-positive cells in the pathogenesis of LN. This may have important implications for therapy, the first open-label pilot study using the senolytic drugs dasatinib plus quercetin (DQ) having shown promising results in idiopathic pulmonary fibrosis and diabetic kidney disease.^{19 20}

Finally, the association of high baseline p16^{INK4a} with impaired renal function 5 years post-treatment initiation suggests it may be a promising predictor of disease severity. It would be of interest to assess for p16^{INK4a} and its associated markers (fibrosis and CD8⁺ T cell infiltration) in 1-year follow-up biopsies as well as in a larger cohort, as the evolution of these markers from baseline to 1 year may better reflect treatment response than either time point alone.

Author affiliations

¹Genetics of Autoimmune Diseases and Cancer, de Duve Institute, Université catholique de Louvain, Brussels, Belgium

²Department of Rheumatology, Cliniques universitaires Saint-Luc, Brussels, Belgium

³IREC Imaging Platform (2IP), Institut de Recherche Expérimentale et Clinique, Université catholique de Louvain, Brussels, Belgium

⁴Department of Pathology, Cliniques Universitaires Saint-Luc, Brussels, Belgium

⁵de Duve Institute, Université catholique de Louvain, Brussels, Belgium

⁶Walloon Excellence in Life Sciences and Biotechnology, Brussels, Belgium

⁷Pôle de pathologies rhumatismales systémiques et inflammatoires, Institut de Recherche Expérimentale et Clinique, Université catholique de Louvain, Brussels, Belgium

Acknowledgements We acknowledge patients and their families, and Prof Anabelle Decottignies for valuable scientific discussions.

Contributors GT, BL and NL: planning, acquisition and analysis of data, drafting of the manuscript; CB: acquisition and analysis of data; SA, FT, CG and FH: acquisition and analysis of clinical data; FH and PGC: planning and drafting of the manuscript.

Funding NL is a chercheur qualifiée of the F.R.S.-F.N.R.S. (Fonds de la Recherche Scientifique, Belgium). This work was supported by the Fonds de la Recherche Fondamentale Stratégique-WELBIO (Walloon Excellence in Life Sciences and Biotechnology), Belgium (Grant WELBIO-CR-2019A-03R) and Actions de Recherche Concertées, UCLouvain (A.R.C. grant 19/24-098), Belgium.

Competing interests Bernard Lauwerys is currently employed at UCB Pharma, Anderlecht, Belgium.

Patient and public involvement statement Patients or the public were not involved in the design, or conduct, or reporting, or dissemination plans of our research.

Patient consent for publication Not applicable.

Ethics approval This study was approved by the Ethics Committee of Cliniques universitaires Saint-Luc (2016/01FEV/034).

Provenance and peer review Not commissioned; externally peer reviewed.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iD

Nisha Limaye <http://orcid.org/0000-0002-9820-4794>

REFERENCES

- Hahn BH. Antibodies to DNA. *N Engl J Med* 1998;338:1359–68.
- Fanouriakis A, Kostopoulou M, Alunno A, et al. 2019 update of the EULAR recommendations for the management of systemic lupus erythematosus. *Ann Rheum Dis* 2019;78:736–45.
- Hanly JG, O'Keefe AG, Su L, et al. The frequency and outcome of lupus nephritis: results from an international inception cohort study. *Rheumatology* 2016;55:252–62.
- Calcinotto A, Kohli J, Zagato E, et al. Cellular senescence: aging, cancer, and injury. *Physiol Rev* 2019;99:1047–78.
- Hernandez-Segura A, Nehme J, Demaria M. Hallmarks of cellular senescence. *Trends Cell Biol* 2018;28:436–53.
- Coppé J-P, Desprez P-Y, Krtolica A, et al. The senescence-associated secretory phenotype: the dark side of tumor suppression. *Annu Rev Pathol* 2010;5:99–118.
- Valentijn FA, Falke LL, Nguyen TQ, et al. Cellular senescence in the aging and diseased kidney. *J Cell Commun Signal* 2018;12:69–82.
- Yang C, Xue J, An N, et al. Accelerated glomerular cell senescence in experimental lupus nephritis. *Med Sci Monit* 2018;24:6882–91.
- Couzi L, Merville P, Deminière C, et al. Predominance of CD8+ T lymphocytes among periglomerular infiltrating cells and link to the prognosis of class III and class IV lupus nephritis. *Arthritis Rheum* 2007;56:2362–70.
- Pamfil C, Makowska Z, De Groof A, et al. Intrarenal activation of adaptive immune effectors is associated with tubular damage and impaired renal function in lupus nephritis. *Ann Rheum Dis* 2018;77:1782–9.
- Braun H, Schmidt BMW, Raiss M, et al. Cellular senescence limits regenerative capacity and allograft survival. *J Am Soc Nephrol* 2012;23:1467–73.
- Baker DJ, Childs BG, Durik M, et al. Naturally occurring p16(Ink4a)-positive cells shorten healthy lifespan. *Nature* 2016;530:184–9.
- Moiseeva O, Mallette FA, Mukhopadhyay UK, et al. Dna damage signaling and p53-dependent senescence after prolonged beta-interferon stimulation. *Mol Biol Cell* 2006;17:1583–92.
- Acosta JC, Banito A, Wuestefeld T, et al. A complex secretory program orchestrated by the inflammasome controls paracrine senescence. *Nat Cell Biol* 2013;15:978–90.
- Gao L, Bird AK, Meednu N, et al. Bone marrow-derived mesenchymal stem cells from patients with systemic lupus erythematosus have a senescence-associated secretory phenotype mediated by a mitochondrial antiviral signaling Protein-Interferon-β feedback loop. *Arthritis Rheumatol* 2017;69:1623–35.
- Lazzeri E, Ronconi E, Angelotti ML, et al. Human urine-derived renal progenitors for personalized modeling of genetic kidney disorders. *J Am Soc Nephrol* 2015;26:1961–74.
- McHugh D, Gil J. Senescence and aging: causes, consequences, and therapeutic avenues. *J Cell Biol* 2018;217:65–77.
- Pereira BI, Devine OP, Vukmanovic-Stejić M, et al. Senescent cells evade immune clearance via HLA-E-mediated NK and CD8+ T cell inhibition. *Nat Commun* 2019;10:2387.
- Hickson LJ, Langhi Prata LGP, Bobart SA, et al. Senolytics decrease senescent cells in humans: preliminary report from a clinical trial of dasatinib plus quercetin in individuals with diabetic kidney disease. *EBioMedicine* 2019;47:446–56.
- Justice JN, Nambiar AM, Tchkonja T, et al. Senolytics in idiopathic pulmonary fibrosis: results from a first-in-human, open-label, pilot study. *EBioMedicine* 2019;40:554–63.