REVIEW

Call for action: incorporating wellness practices into a holistic management plan for rheumatoid arthritis – going beyond treat to target

Peter C. Taylor ¹, Mart Van de Laar,² Andrew Laster,³ Walid Fakhouri,⁴ Amanda Quebe ⁴, Inmaculada de la Torre,⁴ Sandra Jain⁵

To cite: Taylor PC, Van de Laar M, Laster A, *et al.* Call for action: incorporating wellness practices into a holistic management plan for rheumatoid arthritis—going beyond treat to target. *RMD Open* 2021;**7**:e001959. doi:10.1136/rmdopen-2021-001959

Received 22 September 2021
Accepted 23 November 2021

© Author(s) (or their employer(s)) 2021. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

¹Botnar Research Centre, NDORMS, University of Oxford, Oxford, UK

²Transparency in Healthcare B.V., University of Twente, Hengelo, The Netherlands

³Arthritis & Osteoporosis Consultants of the Carolinas, Charlotte, North Carolina, USA

⁴Eli Lilly and Company, Indianapolis, Indiana, USA

⁵School of Nursing, The University of Texas at Austin, Austin, Texas, USA

Correspondence to

Professor Peter C. Taylor;
peter.taylor@kennedy.ox.ac.uk

ABSTRACT

This expert opinion article explores the strategy of adopting a holistic approach to the management of rheumatoid arthritis (RA) by incorporating the wellness practices of exercise, optimised sleep, optimised nutrition, mindfulness, social connectedness and positive emotions into the management plan. The aim is to attain optimal health for each patient beyond that achievable by limiting disease management to pharmacological treatment to attain the lowest achievable composite scores of disease activity, as recommended with the current treat-to-target approach, and addressing the recent recognition of pain control as a key patient-reported outcome. Incorporating wellness practices into a busy clinical setting requires creativity and customisation based on the individual practice setting and the individual needs of each patient. Such practices can help people living with RA to achieve optimum wellness through the introduction of measures—according to individual need—designed to improve the aspects of life most impacted for that person, thereby complementing treat-to-target and pain control strategies with pharmacological agents. Clinicians must consider wellness practices in addition to treat-to-target pharmacological agents for the holistic management of people with RA.

INTRODUCTION

Wellness can be defined as ‘an active process through which people become aware of, and make choices toward, a more successful existence’.¹ It is a multidimensional, holistic concept encompassing lifestyle, environment and mental and spiritual wellbeing. Wellness differs from health promotion (eg, stopping smoking) and comorbidity control (eg, lipid control to prevent heart disease) as it encompasses attitudes and active decisions that contribute to positive health outcomes. Wellness can be improved through a combination of exercise, optimum nutrition, social support, constructive coping strategies (eg, optimised sleep practices and mindfulness) and personal responsibility.¹ Previous

Key messages

- ▶ Wellness practices (exercise, optimised sleep, optimised nutrition, mindfulness, social connectedness and positive emotions) can help people with rheumatoid arthritis (RA) to improve their health status by reducing inflammation and symptoms (eg, stiffness, pain, fatigue) and improving functional ability and wellbeing.
- ▶ Such practices complement and enhance outcomes achievable with treat-to-target pharmacological management alone and encourage a more complete and patient-centred approach to treating RA in individuals with ongoing symptoms.
- ▶ Incorporating wellness practices into a busy clinical setting requires creativity and customisation based on the individual practice setting and the individual needs of each patient.
- ▶ Clinicians should consider wellness practices in addition to treat-to-target pharmacological agents for the holistic management of people with RA.

research highlighted the value of these wellness interventions in chronic diseases, particularly rheumatoid arthritis (RA).²

This expert opinion article explores the science and strategy of a holistic approach to treating RA by incorporating wellness practices into the management plan to help patients achieve optimal health in addition to that achievable with current treat-to-target^{3 4} and patient-reported outcome (PRO) improvement strategies⁵ supported by the use of novel antirheumatic drugs.⁶

METHODOLOGY

This article used references identified through non-systematic searches of the internet, including Medline and Embase, using the search terms ‘rheumatoid arthritis’, ‘exercise’, ‘lifestyle’, ‘diet’, ‘sleep’, ‘wellness’,

Table 1 Exercise and dietary recommendations for patients with rheumatoid arthritis^{10 11}

Exercise	Beneficial effects	Schedule for optimum results
Exercise recommendations		
Walking	Helps with aerobic conditioning and mood	30–60 min, 3–5 times/week
Stretching	Helps with flexibility and range of motion	10–15 min, 2 times/week
Flowing movements	Helps with flexibility, range of motion, balance and stress	10–15 min, 2 times/week
Working out in water	Helps with flexibility, range of motion, aerobic conditioning and strength	30–60 min, 3–5 times/week
Cycling	Improves range of motion, aerobic conditioning, endurance and leg strength	30–60 min, 3–5 times/week
Strength training	Helps with strength and aerobic condition	8–12 repetitions for 2–3 sets, 2–3 times/week
Hand exercises	Improves range of motion and flexibility	
Dietary recommendations—Mediterranean diet		
High intake	Fruit; vegetables; legumes; nuts; unrefined whole grains (whole wheat, rice, oats, corn, barley, rye); unsaturated fats (olive oil, canola/rapeseed oil)	
Moderate intake	Lean meats; fish; wine	
Low-to-moderate intake	Dairy products	
Low intake	Red meat; processed foods; saturated fat	

‘social support’, ‘mindfulness’, ‘positive emotions’, ‘self-care’, and ‘patient-reported outcomes’.

Exercise

Up to 80% of patients with RA have limited exercise capacity due to joint inflammation and damage,⁷ leading to physical pain and functional limitation.⁸ Exercise prevents muscular atrophy and improves physical function by maintaining/improving muscular strength and range of motion, and reducing pain.⁹ Ideally, the patient, rheumatologist and physical therapist should collaborate to ensure exercise undertaken is appropriate for each disease phase to prevent further joint damage. The best exercises are walking, stretching, flowing movements, working-out in water, cycling and strength training^{10 11} (table 1).

Perceived barriers to exercise from the perspective of patients with RA include pain, fatigue, fear of damaging joints, comorbidities, insufficient advice from healthcare providers (HCPs) and lack of time or support.¹² However, regular exercise can reduce disease activity, pain and fatigue and improve physical function, sleep quality and quality of life (QoL) in patients with RA.^{11 13–17} Additionally, it can improve cardiorespiratory/cardiovascular health, muscle mass and strength and reduce adiposity.^{10 13 15}

Possible mechanisms for the beneficial effects of exercise in RA include stimulation of interleukin-6 release from skeletal muscle; increased angiogenesis, leading to reduced hypoxia and associated inflammation; reduced endothelial cell production of adhesion molecules, stimulating regeneration of these cells and reduced vascular wall inflammation; reduced expression of toll-like receptors and proinflammatory cytokine production in monocytes and increased regulatory T-cell production.¹⁸

Sleep

Disturbed sleep is a major concern for people with RA.^{19 20} Many studies report relationships between pain, depression, sleep and functional disability in RA,²¹ including interdependence in causality.²² Sleep and pain should be routinely evaluated during clinical assessments using a multidimensional PRO tool, such as the Rheumatoid Arthritis Impact of Disease Score,²³ and non-pharmacological management approaches, such as physiotherapy, meditation, massage, sleep restriction therapy, sleep scheduling and imagery exercises, advised.²⁴ Practical approaches for enhancing sleep and wellbeing are summarised in box 1.

Poor sleep is one of the risk factors for RA²⁵ and can lead to increased pain, fatigue, depression and anxiety.²⁶ Lack of pain control also impacts sleep, resulting in a vicious cycle.²⁷ Immune activation, as occurs in active RA, can disrupt both deep and dreaming sleep, leading to sleep fragmentation, feeling unrefreshed and daytime fatigue.²⁸ Medications, such as beta-blockers, corticosteroids, analgesics and antidepressants, may further compound the problem.²⁹ Improved sleep through effective RA management can improve disease activity measures.³⁰

Sleep deprivation reduces ATP production and alters the lymphatic system, leading to reduced clearance of cellular waste products. Hyperphosphorylated tau and amyloid β plaques accumulate, increasing cell death and, ultimately, causing cognitive dysfunction.³¹ Persistent sleep disturbance activates β -adrenergic signalling, increasing inflammatory gene expression, proinflammatory cytokine production and markers of systemic inflammation. It also increases monocyte production of signal transducer and activator of transcription proteins, which

Box 1 Techniques to promote effective sleep and improve social connectedness^{77–79}

Promoting effective sleep

- ▶ Exercise regularly.
- ▶ Manage stress through stress-management strategies.
- ▶ Avoid daytime naps.
- ▶ Have a regular sleeping schedule; try to get 7–8 hours of sleep.
- ▶ Do not go to bed unless you are feeling sleepy.
- ▶ Avoid caffeine, alcohol, and nicotine close to bedtime.
- ▶ Do not go to bed hungry, but also do not have a heavy meal close to bedtime.
- ▶ Reduce fluid intake before bedtime.
- ▶ Avoid the use of electronic devices at least 30 min before bedtime.
- ▶ Limit exposure to bright light close to bedtime.
- ▶ Develop a relaxing bedtime routine (eg, have a warm bath, read, listen to relaxing music).
- ▶ Keep the bedroom quiet, cool and dark.

Improving social connectedness

- ▶ Join a volunteer group.
- ▶ Get involved with community activities or a cause the person is passionate about.
- ▶ Take up a hobby and join an appropriate group.
- ▶ Take up a team sport.
- ▶ Keep in contact with family and friends.

mediate inflammatory cytokine signalling, and impacts the sympathetic nervous system and hypothalamic–pituitary–adrenal axis, which are also involved in inflammatory signalling.³²

Nutrition

Of all the wellness behaviours with a potential impact on RA, patients are most likely to ask about diet. Currently, a Mediterranean diet is the most studied dietary recommendation, although other diets have potential anti-inflammatory benefits.³³ Recommended components of the Mediterranean diet are listed in [table 1](#).¹¹

Studies suggest that a Mediterranean (vs non-Mediterranean) diet is associated with a reduced risk of developing RA, particularly in seropositive individuals,^{34 35} and improves inflammation, joint swelling and physical function.^{36 37} The Mediterranean diet derives its benefit from polyunsaturated fatty acids (PUFA), polyphenols and fibre. PUFA, found in oily fish and fish oils, have been shown to reduce inflammatory cytokine levels, while also increasing anti-inflammatory lipid production.³⁸ Polyphenols—a family of phytochemicals in fruits, vegetables and olive oil—reduce oxidative stress and inflammation, while dietary fibre—in whole grains, fruit and vegetables—demonstrates anti-inflammatory properties.³⁹ A Mediterranean diet may reduce the risk of atherosclerotic cardiovascular and cerebrovascular disease, which are increased in patients with active RA.⁴⁰

HCPs should set patient expectations so dietary changes are viewed as complementary rather than an alternative to pharmacological therapy, as the benefits are likely to be modest.³⁸ Of note, dietary effects in patients with

RA may vary with sex, serologic status, gut microbiome and hormonal status (in women).^{33–35} Weight loss discussions, particularly with patients who are obese, can help patients to lose weight⁴¹ and form an important first step in changing patients' weight management behaviour.⁴² Advice on appropriate weight-management services should be provided, while recognising factors that demotivate/motivate patients. Barriers to weight loss include pain and fatigue, limited mobility, stress, work demands and other priorities, poor psychological health, comorbidities and boredom. Appropriate advice includes increased intake of fruit, vegetables and water, switching to low calorie food/drinks, reduced intake of unhealthy foods and drink, reduced portion sizes, changes in eating habits and increased physical activity.⁴³

Mindfulness

Mindfulness can be defined as 'the awareness that arises from paying attention, on purpose, in the present moment and non-judgmentally to the unfolding of experience moment by moment'.⁴⁴ Mindfulness meditation is the wellness practice most likely to be met with uncertainty, probably being a new concept for many HCPs and patients.

To demystify mindfulness, it is helpful to recommend mindfulness apps and provide a one-page handout on its benefits (reduced stress, pain, anxiety and depression and improved health and wellbeing),⁴⁵ along with a list of online free guided meditations and 'how-to' suggestions⁴⁶ ([table 2](#)). Educating patients about how little time is needed to meditate is also helpful; 10 min of mindfulness meditation per day is beneficial, particularly in patients with anxious and repetitive thoughts,⁴⁷ while 20 min/day for 3 days can reduce pain and anxiety.⁴⁸ Tips for fitting meditation into everyday life include introducing the practice into all aspects of daily life (eg, when walking, sitting at one's desk), having a set routine, practicing for short sessions (a few minutes), using reminders (eg, a sound from a watch or computer) and using a meditation anchor (eg, observing the breath).⁴⁹

Mindfulness training can reduce disease activity, physical disability^{17 50} and inflammation⁵¹ and improve morning stiffness, pain, fatigue, patient QoL, illness perception and psychological aspects of RA (eg, depression, anxiety, stress).^{17 50–53} Stress exacerbates RA symptoms, including pain, and can cause immune dysregulation, increased proinflammatory cytokine production, cardiovascular reactivity and altered coagulation.⁵⁴ Mindfulness-based stress reduction significantly increases grey matter concentration in the left hippocampus of the brain—an area involved in learning and memory processes, emotion regulation, self-referential processing and perspective taking⁵⁵—and activates brain regions that modulate pain.⁵⁶

Social connectedness

One in three patients with RA believe others do not understand the impact of their disease and more than

Table 2 Suggestions for practicing mindfulness⁴⁶

Stop and breathe	▶ Take frequent breaks and breathe deeply several times during the day; this fosters calmness and focus
Take time to sit still	▶ Sit quietly in purposeful thought and reflection each day ▶ Avoid filling time with activities
Focus on one task at a time	▶ Practice moment-to-moment awareness in everyday activities
Listen well to everyone	▶ Listen actively and mindfully to coworkers, family members and friends; strong relationships form a strong support network
Appreciate the world around you	▶ Take a walk and use your senses to enjoy what surrounds you
Feed your body well	▶ Choose seasonal, colourful food containing healthy phytonutrients
Eat mindfully	▶ Eat slowly and enjoy the sight, taste and smell of food; it allows for easier digestion
Practice gratitude	▶ Write down five things you are grateful for three times per week
Prepare for bed	▶ Soothe yourself at the end of each day ▶ 1 hour before bedtime, dim the lights, set aside electronic equipment, take a warm bath, read

half feel frustrated when unable to undertake daily activities.⁵⁷ Additionally, many patients believe rheumatologists do not support subjective interventions or consider patients' personal situations as part of the disease control process. However, by suggesting sources of support (eg, community groups), HCPs can help to improve an individual's social connectedness.⁵⁸

Social connections with other people in the same situation and feelings of inclusion are important for physical and mental health.⁵⁹ Support from family and friends is also important to help patients with RA to better understand their illness, attend appointments, adhere to treatments and eat an appropriate diet.⁶⁰ Further suggestions for improving social connections are shown in [box 1](#).

Rewarding social relationships are known to predict better mental and physical health and greater longevity,^{61 62} while loneliness and social isolation are associated with increased inflammation and poor health and wellbeing.^{63 64} Conversely, pain and physical impairment can impact patients' social and work lives.⁶⁵

Social isolation promotes immune dysfunction, including proinflammatory cytokine production, and increases inflammatory markers. Such isolation may cause stress or depression that stimulates proinflammatory cytokine production and exacerbates inflammatory symptoms.^{54 64} Additionally, social isolation may exacerbate the adverse effects of disturbed sleep on inflammation.³²

Positive emotions

Positive emotions (joy, contentment, happiness, love, optimism, serenity and amusement) contribute to mental and physical wellbeing.^{61 66 67} However, HCPs are often trained to focus on negative emotions and reducing symptoms. Integrating positive emotions into clinical work allows HCPs to go beyond this more traditional treatment approach. Encouraging patients to undertake an activity they enjoy is similar to prescribing exercise, social connectedness or other wellness practices; takes little time and is cost effective. Similar practices can be

individualised and integrated into treatment planning and may even improve RA medication adherence.⁶⁸

The link between positive emotions and physical health and longevity is well known. Positive emotions also promote positive social relationships that are reciprocally associated with physical health,⁶¹ reduce the intensity of persistent pain and buffer the effects of such pain on functioning and wellbeing in patients with RA.⁶⁹

Studies have shown that positive emotions are associated with larger grey matter volume in a brain area from the left thalamus to the parahippocampal gyrus⁷⁰ and activate brain areas involved in the sensing of pleasure (ventral striatum and orbital frontal cortex).⁷¹ Positive emotions are also associated with lower circulating proinflammatory cytokine levels.^{66 67 72}

DISCUSSION AND 'CALL FOR ACTION'

Treat to target has been the main approach to managing RA, based on compelling evidence for reduced joint destruction and preservation of functional status. However, many areas of contemporary unmet need are subjective aspects of key importance to the individual, such as pain reduction, improved functioning and QoL. While pharmacological interventions are needed, it is important to provide a holistic approach to treating patients to reduce pressure on healthcare systems and increase productivity. Studies have highlighted the importance of the emotional and social impact of RA and how most patients still do not feel well despite pharmacological treatment, mainly due to uncontrolled pain that impacts daily activities and wellbeing.⁵⁷

As outlined in this article, wellness interventions can improve health and reduce the impact of RA on a person from both a physical and an emotional perspective. These lifestyle changes can be considered part of the overarching practice of self-management, which encompasses a variety of activities aimed at improving a person's ability to manage symptoms and treatment, and the physical and psychological consequences of a

chronic disease.⁷³ The importance of self-management strategies to optimise wellbeing for people living with RA has recently been recognised with the publication of EULAR recommendations⁷⁴ and Portuguese multi-disciplinary recommendations for non-pharmacological interventions.⁷⁵ The latter include exercise, psychological interventions, social participation, sleep hygiene and self-management.⁷⁵ Recommended self-management strategies include problem solving, goal setting and cognitive behavioural therapy (CBT).⁷⁴ CBT is a type of psychotherapy used to challenge negative thought patterns to change patient behaviours that maintain symptoms and disability; it has been shown to provide small-to-moderate improvements in impairment/disability, fatigue and self-efficacy in patients with RA.^{17 76}

This is the first article to comprehensively discuss a rationale and existing evidence for the holistic management of RA and to provide practical suggestions for incorporating wellness practices into a busy clinical setting—a process requiring creativity and customisation based on individual needs and practice settings in line with EULAR treat-to-target and self-management goals.^{4 74} Wellness is not necessarily an additional outcome to address above and beyond treat to target, as all treat-to-target metrics include the Patient Global Assessment, which is a PRO. By enhancing wellness and thereby improving patient global scores, rheumatologists may be able to improve other disease metrics (eg, the Disease Activity Score for 28-joint count and the Clinical Disease Activity Index). Whether specific RA drugs also contribute to enhanced wellness by improving key PROs connected to wellbeing (eg, pain) is not yet determined.

We call for clinicians to consider wellness practices in addition to pharmacological agents and PRO measurement for the holistic management of people with RA.

Acknowledgements The authors would like to acknowledge Dr Sue Chambers, Dr Ioannis Nikas and Karen Goa (Rx Communications, Mold, UK) for medical writing assistance with the preparation of this article, funded by Eli Lilly and Company. PT would like to thank the National Institute of Health Research (NIHR) for their funding of the NIHR Biomedical Research Centre in Musculoskeletal Disease at Oxford University Hospitals NHS Trust and University of Oxford.

Contributors Conception of the work: PT, MVDL, IdIT, WF and AQ. Design of the work: PT, IdIT, WF and AQ. Acquisition of data for the work and analysis of data for the work: PT, SJ, AL, IdIT, WF and AQ. Interpretation of data for the work, drafting of the article, critical revision of the article for important intellectual content, approval and sufficient participation in the work to agree to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved: all authors.

Funding This article was funded by Eli Lilly and Company.

Competing interests PT: Research grants from Celgene and Galapagos and consultation fees from AbbVie, Biogen, Galapagos, Gilead, GlaxoSmithKline, Janssen, Lilly, Pfizer, BMS, Roche, Sanofi, Nordic Pharma, Fresenius and UCB. SJ has served as a consultant, member of advisory boards and/or speaker bureaus for Eli Lilly, Otsuka, Pamlab and Sunovion. MVDL reports educational grants and speaker's-advisory fees from Eli Lilly and Company not related to this work. AL has been member of speaker bureaus for Lilly, Novartis, Amgen, Exage, Myria and Pfizer and a consultant for Amgen. WF, AQ and IdIT are employees of Eli Lilly and Company.

Patient consent for publication Not applicable.

Ethics approval This study does not involve human participants.

Provenance and peer review Not commissioned; externally peer reviewed.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Peter C. Taylor <http://orcid.org/0000-0001-7766-6167>

Amanda Quebec <http://orcid.org/0000-0001-6705-8045>

REFERENCES

- 1 National Wellness Institute. The six dimensions of wellness. Available: <https://nationalwellness.org/resources/six-dimensions-of-wellness/> [Accessed 12 Oct 2020].
- 2 Stuifbergen AK, Morris M, Jung JH, *et al.* Benefits of wellness interventions for persons with chronic and disabling conditions: a review of the evidence. *Disabil Health J* 2010;3:133–45.
- 3 Fraenkel L, Bathon JM, England BR, *et al.* American College of Rheumatology guideline for the treatment of rheumatoid arthritis. *Arthritis Rheumatol* 2021;2021:1108–23.
- 4 Smolen JS, Landewé RBM, Bijlsma JWW, *et al.* EULAR recommendations for the management of rheumatoid arthritis with synthetic and biological disease-modifying antirheumatic drugs: 2019 update. *Ann Rheum Dis* 2020;79:685–99.
- 5 Fautrel B, Alten R, Kirkham B, *et al.* Call for action: how to improve use of patient-reported outcomes to guide clinical decision making in rheumatoid arthritis. *Rheumatol Int* 2018;38:935–47.
- 6 Fautrel B, Zhu B, Taylor PC, *et al.* Comparative effectiveness of improvement in pain and physical function for baricitinib versus adalimumab, tocilizumab and tofacitinib monotherapies in rheumatoid arthritis patients who are naïve to treatment with biologic or conventional synthetic disease-modifying antirheumatic drugs: a matching-adjusted indirect comparison. *RMD Open* 2020;6:e001131.
- 7 Sokka T, Häkkinen A. Poor physical fitness and performance as predictors of mortality in normal populations and patients with rheumatic and other diseases. *Clin Exp Rheumatol* 2008;26:S14–20.
- 8 Smolen JS, Aletaha D, McInnes IB. Rheumatoid arthritis. *Lancet* 2016;388:2023–38.
- 9 Hurkmans E, van der Giesen FJ, Vliet Vlieland TP, *et al.* Dynamic exercise programs (aerobic capacity and/or muscle strength training) in patients with rheumatoid arthritis. *Cochrane Database Syst Rev* 2009;2009:CD006853.
- 10 Cooney JK, Law R-J, Matschke V, *et al.* Benefits of exercise in rheumatoid arthritis. *J Aging Res* 2011;2011:681640.
- 11 Saulle R, La Torre G. The Mediterranean diet, recognized by UNESCO as a cultural heritage of humanity. *Ital J Public Health* 2010;7:414–5.
- 12 Veldhuijzen van Zanten JJCS, Rouse PC, Hale ED, *et al.* Perceived barriers, facilitators and benefits for regular physical activity and exercise in patients with rheumatoid arthritis: a review of the literature. *Sports Med* 2015;45:1401–12.
- 13 de Jong Z, Munneke M, Kroon HM, *et al.* Long-term follow-up of a high-intensity exercise program in patients with rheumatoid arthritis. *Clin Rheumatol* 2009;28:663–71.
- 14 Durcan L, Wilson F, Cunnane G. The effect of exercise on sleep and fatigue in rheumatoid arthritis: a randomized controlled study. *J Rheumatol* 2014;41:1966–73.
- 15 Manning VL, Hurley MV, Scott DL, *et al.* Education, self-management, and upper extremity exercise training in people with rheumatoid arthritis: a randomized controlled trial. *Arthritis Care Res* 2014;66:217–27.
- 16 Ambrosino P, Iannuzzi GL, Formisano R, *et al.* Exergaming as an additional tool in rehabilitation of young patients with rheumatoid arthritis: a pilot randomized controlled trial. *Games Health J* 2020;9:368–75.
- 17 Marques A, Santos E, Nikiphorou E, *et al.* Effectiveness of self-management interventions in inflammatory arthritis: a systematic review informing the 2021 EULAR recommendations for the implementation of self-management strategies in patients with inflammatory arthritis. *RMD Open* 2021;7:e001647.
- 18 You T, Arsenis NC, Disanzo BL, *et al.* Effects of exercise training on chronic inflammation in obesity: current evidence and potential mechanisms. *Sports Med* 2013;43:243–56.
- 19 Drewes AM, Nielsen KD, Hansen B, *et al.* A longitudinal study of clinical symptoms and sleep parameters in rheumatoid arthritis. *Rheumatology* 2000;39:1287–9.

- 20 Kirwan J, Heiberg T, Hewlett S, *et al.* Outcomes from the patient perspective workshop at OMERACT 6. *J Rheumatol* 2003;30:868–72.
- 21 Luyster FS, Chasens ER, Wasko MCM, *et al.* Sleep quality and functional disability in patients with rheumatoid arthritis. *J Clin Sleep Med* 2011;7:49–55.
- 22 Wolfe F, Michaud K, Li T. Sleep disturbance in patients with rheumatoid arthritis: evaluation by medical outcomes study and visual analog sleep scales. *J Rheumatol* 2006;33:1942–51.
- 23 Gossec L, Dougados M, Rinccheval N, *et al.* Elaboration of the preliminary Rheumatoid Arthritis Impact of Disease (RAID) score: a EULAR initiative. *Ann Rheum Dis* 2009;68:1680–5.
- 24 Tang NKY, Lereya ST, Boulton H, *et al.* Nonpharmacological treatments of insomnia for long-term painful conditions: a systematic review and meta-analysis of patient-reported outcomes in randomized controlled trials. *Sleep* 2015;38:1751–64.
- 25 Sivertsen B, Lallukka T, Salo P, *et al.* Insomnia as a risk factor for ill health: results from the large population-based prospective HUNT study in Norway. *J Sleep Res* 2014;23:124–32.
- 26 Irwin MR, Olmstead R, Carrillo C, *et al.* Sleep loss exacerbates fatigue, depression, and pain in rheumatoid arthritis. *Sleep* 2012;35:537–43.
- 27 Grabovac I, Haider S, Berner C, *et al.* Sleep quality in patients with rheumatoid arthritis and associations with pain, disability, disease duration, and activity. *J Clin Med* 2018;7:336.
- 28 Besedovsky L, Lange T, Haack M. The sleep-immune crosstalk in health and disease. *Physiol Rev* 2019;99:1325–80.
- 29 Van Gastel A. Drug-Induced insomnia and excessive sleepiness. *Sleep Med Clin* 2018;13:147–59.
- 30 Wells G, Li T, Tugwell P. Investigation into the impact of abatacept on sleep quality in patients with rheumatoid arthritis, and the validity of the MOS-sleep questionnaire sleep disturbance scale. *Ann Rheum Dis* 2010;69:1768–73.
- 31 Lucke-Wold BP, Smith KE, Nguyen L, *et al.* Sleep disruption and the sequelae associated with traumatic brain injury. *Neurosci Biobehav Rev* 2015;55:68–77.
- 32 Irwin MR, Opp MR. Sleep health: reciprocal regulation of sleep and innate immunity. *Neuropsychopharmacol Rev* 2017;42:129–55.
- 33 Coras R, Murillo-Saich JD, Guma M. Circulating pro- and anti-inflammatory metabolites and its potential role in rheumatoid arthritis pathogenesis. *Cells* 2020;9:827.
- 34 Hu Y, Sparks JA, Malspeis S, *et al.* Long-term dietary quality and risk of developing rheumatoid arthritis in women. *Ann Rheum Dis* 2017;76:1357–64.
- 35 Johansson K, Askling J, Alfredsson L, *et al.* Mediterranean diet and risk of rheumatoid arthritis: a population-based case-control study. *Arthritis Res Ther* 2018;20:175.
- 36 Sködstam L, Hagfors L, Johansson G. An experimental study of a Mediterranean diet intervention for patients with rheumatoid arthritis. *Ann Rheum Dis* 2003;62:208–14.
- 37 Vadell AKE, Bärebring L, Hulander E, *et al.* Anti-inflammatory diet in rheumatoid arthritis (ADIRA)-a randomized, controlled crossover trial indicating effects on disease activity. *Am J Clin Nutr* 2020;111:1203–13.
- 38 Miles EA, Calder PC. Influence of marine n-3 polyunsaturated fatty acids on immune function and a systematic review of their effects on clinical outcomes in rheumatoid arthritis. *Br J Nutr* 2012;107:S171–84.
- 39 Bustamante MF, Agustín-Pérez M, Cedola F, *et al.* Design of an anti-inflammatory diet (ITIS diet) for patients with rheumatoid arthritis. *Contemp Clin Trials Commun* 2020;17:100524.
- 40 Crowson CS, Liao KP, Davis JM, *et al.* Rheumatoid arthritis and cardiovascular disease. *Am Heart J* 2013;166:622–8.
- 41 Pool AC, Kraschnewski JL, Cover LA, *et al.* The impact of physician weight discussion on weight loss in US adults. *Obes Res Clin Pract* 2014;8:e131–9.
- 42 Halbert CH, Jefferson M, Melvin CL, *et al.* Provider advice about weight loss in a primary care sample of obese and overweight patients. *J Prim Care Community Health* 2017;8:239–46.
- 43 Evans EH, Sainsbury K, Kwasnicka D, *et al.* Support needs of patients with obesity in primary care: a practice-list survey. *BMC Fam Pract* 2018;19:6.
- 44 Kabat-Zinn J. Mindfulness-based interventions in context: past, present, and future. *Clin Psychol Sci Pract* 2003;10:144–56.
- 45 Tang Y-Y, Leve LD. A translational neuroscience perspective on mindfulness meditation as a prevention strategy. *Transl Behav Med* 2016;6:63–72.
- 46 Cleveland Clinic. Practice mindfulness to improve your well-being: 11 tips, 2018. Available: <https://health.clevelandclinic.org/practice-mindfulness-to-improve-your-well-being-11-tips/> [Accessed 8 Dec 2020].
- 47 Xu M, Purdon C, Seli P, *et al.* Mindfulness and mind wandering: the protective effects of brief meditation in anxious individuals. *Conscious Cogn* 2017;51:157–65.
- 48 Zeidan F, Gordon NS, Merchant J, *et al.* The effects of brief mindfulness meditation training on experimentally induced pain. *J Pain* 2010;11:199–209.
- 49 Shonin E, Van Gordon W, Griffiths MD. Practical tips for using mindfulness in general practice. *Br J Gen Pract* 2014;64:368–9.
- 50 El Miedany Y, El Gaafary M, El Arousy N, *et al.* Arthritis education: the integration of patient-reported outcome measures and patient self-management. *Clin Exp Rheumatol* 2012;30:899–904.
- 51 Fogarty FA, Booth RJ, Gamble GD, *et al.* The effect of mindfulness-based stress reduction on disease activity in people with rheumatoid arthritis: a randomised controlled trial. *Ann Rheum Dis* 2015;74:472–4.
- 52 Zangi HA, Mowinckel P, Finset A, *et al.* A mindfulness-based group intervention to reduce psychological distress and fatigue in patients with inflammatory rheumatic joint diseases: a randomised controlled trial. *Ann Rheum Dis* 2012;71:911–7.
- 53 Dalili Z, Bayazi MH. The effectiveness of mindfulness-based cognitive therapy on the illness perception and psychological symptoms in patients with rheumatoid arthritis. *Complement Ther Clin Pract* 2019;34:139–44.
- 54 Liu Y-Z, Wang Y-X, Jiang C-L. Inflammation: the common pathway of stress-related diseases. *Front Hum Neurosci* 2017;11:316.
- 55 Hölzel BK, Carmody J, Vangel M, *et al.* Mindfulness practice leads to increases in regional brain gray matter density. *Psychiatry Res* 2011;191:36–43.
- 56 Zeidan F, Emerson NM, Farris SR, *et al.* Mindfulness meditation-based pain relief employs different neural mechanisms than placebo and sham mindfulness meditation-induced analgesia. *J Neurosci* 2015;35:15307–25.
- 57 Alten R, van de Laar M, De Leonardi F, *et al.* Physical and emotional burden of rheumatoid arthritis: data from RA matters, a web-based survey of patients and healthcare professionals. *Rheumatol Ther* 2019;6:587–97.
- 58 Mossabir R, Morris R, Kennedy A, *et al.* A scoping review to understand the effectiveness of linking schemes from healthcare providers to community resources to improve the health and well-being of people with long-term conditions. *Health Soc Care Community* 2015;23:467–84.
- 59 My Health My Community. Social connection and health, 2019. Available: https://myhealthmycommunity.org/wp-content/uploads/2019/05/MHMC_SocialConnections_web.pdf [Accessed 9 Dec 2020].
- 60 Ginsberg S. Caregivers are vital to arthritis management. Rheumatology Network, 2017. Available: <https://www.rheumatologynetwork.com/view/caregivers-are-vital-arthritis-management> [Accessed 24 Jun 2021].
- 61 Kok BE, Coffey KA, Cohn MA, *et al.* How positive emotions build physical health: perceived positive social connections account for the upward spiral between positive emotions and vagal tone. *Psychol Sci* 2013;24:1123–32.
- 62 Pillemer S, Holtzer R, Blumen HM. Functional connectivity associated with social networks in older adults: a resting-state fMRI study. *Soc Neurosci* 2017;12:242–52.
- 63 Hawkey LC, Cacioppo JT. Loneliness matters: a theoretical and empirical review of consequences and mechanisms. *Ann Behav Med* 2010;40:218–27.
- 64 Yang YC, Li T, Frenk SM. Social network ties and inflammation in U.S. adults with cancer. *Biodemography Soc Biol* 2014;60:21–37.
- 65 Schneider M, Manabile E, Tikly M. Social aspects of living with rheumatoid arthritis: a qualitative descriptive study in Soweto, South Africa - a low resource context. *Health Qual Life Outcomes* 2008;6:54.
- 66 Stellar JE, John-Henderson N, Anderson CL, *et al.* Positive affect and markers of inflammation: discrete positive emotions predict lower levels of inflammatory cytokines. *Emotion* 2015;15:129–33.
- 67 Moreno PI, Moskowitz AL, Ganz PA, *et al.* Positive affect and inflammatory activity in breast cancer survivors: examining the role of affective arousal. *Psychosom Med* 2016;78:532–41.
- 68 Rodrigues JR, Faria DS, Neves JS, *et al.* Positive affect as a predictor of adherence in patients with rheumatoid arthritis. *Acta Rheumatol Port* 2019;44:132–7.
- 69 Hanssen MM, Peters ML, Boselie JJ, *et al.* Can positive affect attenuate (persistent) pain? State of the art and clinical implications. *Curr Rheumatol Rep* 2017;19:80.
- 70 Yang J, Wei D, Wang K, *et al.* Gray matter correlates of dispositional optimism: a voxel-based morphometry study. *Neurosci Lett* 2013;553:201–5.

- 71 Burgdorf J, Panksepp J. The neurobiology of positive emotions. *Neurosci Biobehav Rev* 2006;30:173–87.
- 72 Matsunaga M, Isowa T, Yamakawa K, *et al.* Association between perceived happiness levels and peripheral circulating pro-inflammatory cytokine levels in middle-aged adults in Japan. *Neuro Endocrinol Lett* 2011;32:458–63.
- 73 Zuidema RM, Repping-Wuts H, Evers AWM, *et al.* What do we know about rheumatoid arthritis patients' support needs for self-management? A scoping review. *Int J Nurs Stud* 2015;52:1617–24.
- 74 Nikiphorou E, Santos EJJ, Marques A, *et al.* EULAR recommendations for the implementation of self-management strategies in patients with inflammatory arthritis. *Ann Rheum Dis* 2021;2021:1278–85.
- 75 Santos EJJ, Duarte C, Ferreira RJO, *et al.* Portuguese multidisciplinary recommendations for non-pharmacological and non-surgical interventions in patients with rheumatoid arthritis. *Acta Reumatol Port* 2021;46:40–54.
- 76 Hewlett S, Ambler N, Almeida C, *et al.* Self-management of fatigue in rheumatoid arthritis: a randomised controlled trial of group cognitive-behavioural therapy. *Ann Rheum Dis* 2011;70:1060–7.
- 77 Martino J, Pegg J, Frates EP. The connection prescription: using the power of social interactions and the deep desire for connectedness to empower health and wellness. *Am J Lifestyle Med* 2015;11:466–75.
- 78 Irish LA, Kline CE, Gunn HE, *et al.* The role of sleep hygiene in promoting public health: a review of empirical evidence. *Sleep Med Rev* 2015;22:23–36.
- 79 Sleep Education. Healthy sleep habits, 2020. Available: <https://sleepeducation.org/healthy-sleep/healthy-sleep-habits/> [Accessed 24 Jun 2021].