

ORIGINAL RESEARCH

Impact of synovial biopsy procedures and disease-specific aspects on synovial tissue outcome: a systematic literature review informing the EULAR points to consider for the minimal reporting requirements in synovial tissue research in rheumatology

Aur lie Najm ¹, F licie Costantino ^{2,3}, Catherine Weill,⁴ Andrew Filer ⁵, Maria Antonietta D'Agostino^{2,6}

To cite: Najm A, Costantino F, Weill C, *et al*. Impact of synovial biopsy procedures and disease-specific aspects on synovial tissue outcome: a systematic literature review informing the EULAR points to consider for the minimal reporting requirements in synovial tissue research in rheumatology. *RMD Open* 2022;**8**:e002116. doi:10.1136/rmdopen-2021-002116

► Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/rmdopen-2021-002116>).

Received 18 November 2021

Accepted 15 January 2022



  Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Aur lie Najm;
aurelie.najm@glasgow.ac.uk

ABSTRACT

Background The aim of this work was to summarise the literature evaluating the impact of biopsy procedures, tissue handling, tissue quality and disease-specific aspects including joint biopsied and disease stage, on synovial tissue outcome.

Methods Two reviewers independently identified eligible studies according to the Patients, Intervention, Comparator and Outcome framework obtained for five research questions formulated during the first EULAR task force meeting to produce points to consider (PtC) for minimal reporting requirements in synovial tissue studies. The databases explored were Medline, Embase, CENTRAL and Cinhal. The risk of bias of each study was evaluated using an adapted version of the Joanna Briggs Institute checklist for analytical cross-sectional studies.

Results Of the 7654 records yielded, 75 full texts were assessed, leading to the inclusion of 26 manuscripts in the systematic literature review (SLR). Two papers assessed the impact of biopsy procedures on the quality and quantity of tissue retrieved alongside patient tolerability; six papers focused on synovial tissue variability. Four papers studied the impact of sample handling or randomisation and 14 assessed the impact of disease stage and state, namely early or established active rheumatoid arthritis and remission on histopathological and transcriptomic results.

Conclusions This SLR informs the EULAR PtC for minimal reporting requirements in synovial tissue research in rheumatology. Characteristics related to the study design, population, sample handling, randomisation and analysis can affect the final synovial tissue outcome in the studies reviewed. Thus, accurate reporting of these factors is required in order to ensure the scientific validity of manuscripts describing synovial tissue outcomes.

Key messages

What is already known about this subject?

- Synovial tissue biopsy involves different procedures: ultrasound-guided portal and forceps, ultrasound-guided needle biopsy and arthroscopic biopsies.
- While different sampling, handling and analysis techniques are reported in publications, their impact on study outcomes remain unknown.

What does this study add?

- Histological, immunopathological and transcriptomic outcomes of synovial tissue vary across disease stage and activity.
- Different synovial biopsy techniques, provided they involve guidance, do not substantially affect tissue quantity or quality and patient tolerance.
- Intra-articular and interarticular variability appears minimal for histopathological or transcriptomic outcomes in rheumatoid arthritis, if number of sample fragments taken from each joint is adequate.

How might this impact on clinical practice?

- Information related to tissue retrieval, handling, quality and analysis can affect synovial tissue outcome and therefore should be mandatorily reported in scientific manuscripts.

INTRODUCTION

Studies involving synovial tissue (ST) in rheumatic and musculoskeletal diseases (RMDs) have underpinned major breakthroughs in the field of immunology over the past years.^{1–5} This, in addition to the development of mini-invasive biopsy techniques, has led to a significant increase in synovial

biopsy (SB) procedures across centres in Europe and beyond.^{6–11} While SB procedures are becoming widespread, high standards in both retrieving and analysing ST are required in order to allow a robust evolution of the field. An unmet need exists for evidence and consensus-based points to consider (PtC) defining minimum reporting requirements that could ensure interpretability of the research. In this context, the European alliance of associations for rheumatology (EULAR) has approved the constitution of a task force to develop PtC for minimal reporting requirements in ST research in RMDs.

Although it is assumed that several aspects related to ST handling in research can impact the scientific outcomes, the published studies assessing the influence of tissue retrieving, handling and analysis methods on the results remain scarce and sometimes contradictory. In this context, it was felt important to summarise the existing literature describing how aspects of study design and methods can impact the results. In addition, it is unclear how the joint selected for biopsy, the area within the joint where the tissue is retrieved or how samples are randomised to each analysis, potentially affect the outcome of the research. Therefore, the aim of this systematic literature review (SLR) was to identify the impact of biopsy procedures, tissue handling, tissue quality and disease-specific aspects, on tissue outcomes, to inform the EULAR PtC for minimal reporting requirements in ST research in RMDs.

METHODS

Research questions

The SLR was conducted following the EULAR standardised operating procedures.¹² The scope of the literature search was defined during a first virtual task force meeting, in which five different research questions (RQ) were formulated and approved by all task force members and focused on biopsy techniques (RQ1), representativity of ST from large and small joints in inflammatory RMDs (RQ2), sample randomisation and tissue handling (RQ3), tissue quality control (RQ4) and impact of disease stage and activity on tissue outcome (RQ5).

Search methodology

The five RQs were transformed using the 'Patients, Intervention, Comparator and Outcome' (PICO) framework to determine the search strategy (online supplemental text S1).¹³ The search was run by an experienced librarian from Paris University, Paris, France (CW). The keywords selected and used for each PICO and databases are presented as supplementary material (online supplemental text S2). The databases explored were Medline, Embase, CENTRAL and Cinhal. Hand search for individual original research studies and crosscheck for references from specific rheumatology and immunology journals were performed.

Study selection, data extraction and risk of bias assessment

Titles and abstracts of the retrieved papers were assessed by two independent reviewers (AN and FC). General eligibility criteria were described as follows: original research articles, published in peer-reviewed journals, English language, reporting clinical or translational research involving ST. Cohen's kappa agreement between reviewers was 0.95. Discrepancies were resolved by discussion.

Data on patients' characteristics, scientific methods of analysis, parameters assessed and outcomes were extracted. In addition, all manuscripts included in the final SLR were assessed from their content against criteria compelled during the first task force meeting by the task force members. These criteria identified important areas where data should be reported on a mandatory basis.

Due to the lack of validated risk of bias (RoB) tool for translational research in rheumatology, the RoB was assessed using an adapted version of the Joanna Briggs Institute checklist for analytical cross-sectional studies.¹⁴ Briefly, eight questions related to specific methodological aspects were applied to the study. If ≥ 2 'no' answers were returned, the study was considered at high RoB, while RoB was said to be 'unclear' if one 'no' answer was returned. The tool is detailed in online supplemental text S3.

RESULTS

Study selection and study characteristics

Overall, our search yielded 7654 abstracts out of which 2181 duplicates were excluded; 5473 titles and abstracts were screened, and 75 full texts assessed, leading to the inclusion of 26 manuscripts in the SLR. Additional details are presented in the flow chart (figure 1). Study characteristics with their type, objective, number and characteristics of patients and studied outcomes are presented in table 1. The RoB assessment is presented in table 2. Most studies were found to have an unclear (n=14) or high (n=12) RoB mainly due to the existence of confounding factors.

Reporting methods and outcomes across included studies

We additionally analysed the way each included manuscript reported their materials and methods section, with a focus on several areas deemed important by the task force, as listed in table 3. Essentially, manuscripts have been checked for the following criteria: clinical data, biopsy procedure, tissue handling, tissue quality control and tissue outcomes. The area most likely to be reported in all manuscripts were: demographics (100%), disease duration (92%), target joints for biopsy (81%), number of fragments retrieved and included for analysis (73%) and their processing (100%), scoring/quantification system used for immunohistochemistry (IHC, 96%) and disaggregation method used for single-cell analysis (100%). The least likely items to be reported (0%–40%) were: ultrasound (US) features of the biopsied joints




Figure 1 Flow chart summary of the systematic literature review, article identification, screening and final selection. PICO, patient, intervention, comparison, outcome.

(for US-guided procedures), operator's experience and adverse events related to the biopsy procedure, intraobserver or interobserver reproducibility for IHC and RNA quantity, purity and quality control method.

Impact of the biopsy procedure and device on tissue quality and biopsy tolerability

One study with unclear RoB assessed the impact of biopsy procedure and devices on ST yield and quality. In this study, the three main guided biopsy techniques: US-guided needle biopsy (NB), ultrasoundUS-guided portal and forceps (P&F), arthroscopy were compared with each other and with blind NB. The number of graded ST fragments/total number of ST fragments was higher in guided techniques compared with blind NB for both small and large joints although this result was only significant for large joints ($p=0.048$ and $p=0.057$, respectively).¹⁵ While all guided techniques enabled retrieval of a sufficient amount and quality of tissue for a meaningful analysis, large joint arthroscopic-guided biopsy (A) yielded higher quantities of RNA than US-guided NB and US P&F biopsy (A vs NB $p=0.002$, A vs P&F $p=0.0014$) and higher RIN for A compared with NB ($p=0.0018$) but not compared with P&F ($p=0.068$). No difference was observed in immune cell infiltrate and pathotype

regardless of the technique used. For sequential biopsies performed with any of the guided techniques, no difference was observed between first and second biopsies for tissue yield and quality or RNA quality.

As far as tolerance is concerned, one study with unclear RoB assessed the influence of the biopsy technique on patient-reported outcomes.¹⁶ Visual analogic scale (VAS) for pain, swelling and stiffness did not differ for A, NB or P&F, while the willingness to repeat was superior in P&F compared with both A ($p<0.01$) and NB ($p<0.01$).¹⁶ In addition, no difference was observed in terms of reported PROs for tolerance or adverse events between first and second biopsies although numbers were low.

These results are summarised in tables 4 and 5 (sequential biopsies).

Impact of the joint biopsied on tissue outcomes

Two studies with high RoB^{17,18} and one with unclear RoB¹⁹ assessed the impact of the joint biopsied on either cell infiltrate,^{17,19} transcriptomics¹⁹ or T-cell repertoire (TCR), deemed to be particularly relevant in ST arthritis of autoimmune mechanisms.^{18,19} Of interest, when comparing outcomes from tissues retrieved in small versus large joints from the same patients with rheumatoid arthritis (RA), the authors

Table 1 Design, outcomes and population of studies included in the manuscript

RQ	Author and year	Objective	Patients	Procedures	Outcomes
RQ 1	Humby <i>et al</i> 2018 ¹⁵	Comparing across techniques: ▶ ST yield/ST quantity ▶ ST quality	125 patients: ▶ RA (n=73) ▶ UA (n=31) ▶ PsA/SpA (n=18) ▶ JIA (n=3) Follow-up (n=34)	Synovial biopsies (n=159) ▶ US-NB (n=47) ▶ US-P&F (n=54) ▶ Arthroscopy (n=35) ▶ Blind NB (n=23)	Graded ST fragments/total ST fragments Total mean graded ST area Mean ST area per graded biopsy fragment Total mean graded ST surface Mean synovitis scores Mean semiquantitative scores of synovial infiltration in sublining (SL) of CD20 +B cells; CD3 +T cells; CD68 +macrophages Frequency of synovial pathology RNA quantity and RIN in small and large joints
	Just <i>et al</i> 2018 ¹⁶	Comparing across techniques: ▶ Tolerance ▶ Adverse events	421 patients ▶ RA (n=360) ▶ UA (n=20) ▶ PsA/SpA (n=14) ▶ Others (n=8) Follow-up (n=103)	Synovial biopsies (n=524) ▶ US-NB (n=402) ▶ US-P&F (n=65) ▶ Arthroscopy (n=57)	VAS: joint pain/joint swelling/stiffness Delta VAS: joint pain/joint swelling/stiffness 'How does joint feel compared with before biopsy' Willingness to repeat: 1–5 Likert scale Adverse events
RQ2	Kraan <i>et al</i> 2002 ¹⁷	Comparing in paired biopsies from small and large joints: ▶ Cell infiltrate ▶ IL-6 expression	9 RA patients	Arthroscopic SB (n=18) ▶ Knee (n=9) ▶ Paired small joints (MCP: n=4; wrist: n=5)	Cell numbers per mm ² in lining layer (LL) and SL: ▶ CD55 +fibroblast like synoviocytes ▶ CD68 +macrophages ▶ CD3 +T cells ▶ CD138 +plasma cells IL-6 expression (integrated optical density)
	Musters <i>et al</i> 2018 ¹⁸	Comparing in paired biopsies from right and left joints: ▶ T-cell repertoire	9 RA patients	Arthroscopic SB (n=18) ▶ Paired left and right joints (ankle or knee)	Number of TCRβ clones Number of highly expanded clones (HEC) Impact of the HECs on the total repertoire Overlap between the 25 most expanded TCRβ clones (%) Chao-modified Sørensen index
	Triaille <i>et al</i> 2020 ¹⁹	Comparing in paired biopsies from small and large joints: ▶ T-cell repertoire ▶ T-cell signature	10 RA patients	Synovial biopsies (n=20) ▶ Knee (n=10; arthroscopic SB) ▶ Paired small joints (n=10; MCP or wrist; US-NB)	Synovitis scores Mean semiquantitative scores of synovial infiltration of CD20 +B cells, CD3 +T cells and CD68 +macrophages Number of transcripts overexpressed Pathway expression patterns Enriched T-cell clonotypes (ETC) and % of shared ETC

Continued

Table 1 Continued

RQ	Author and year	Objective	Patients	Procedures	Outcomes
RQ3	Donlin <i>et al</i> 2018 ²⁰	Comparing tissue handling methods: ▲ Fresh vs cryopreserved tissue ▲ Dissociation method	RA patients	Synovial biopsies (n=19) ▲ US-NB (n=16) ▲ Surgical biopsies (n=3)	Total cell counts per gram from ST mechanically disaggregated Flow cytometry: ▲ % of total CD90+, CD14+, CD3+, CD19 +cells ▲ % viability
	Smeets <i>et al</i> 2003 ²¹	Comparing intra-articular variability of: ▲ Cell infiltrate ▲ MMPs	17 RA patients	Arthroscopic biopsies (n=17) Paired SB of 2 different sites of the knee (CPJ and SP recess)	Quantitative analysis of CD3, CD4, CD8, CD22, CD38, CD68, MMP-1 and granzyme B7 by computer-assisted image analysis. Semi quantitative analysis of CD3, CD4, CD8, CD22, CD38, CD68, CD55, mast cells, MMP-1, MMP-3 and MMP-13
	Kirkham <i>et al</i> 1999 ²²	Comparing intra-articular variability of: ▲ Cell infiltrate ▲ Cytokines expression	11 RA patients	Arthroscopic biopsies (n=11) Knee; 4 anatomically different sites: CPJ and SP, lateral and medial recesses	Semi quantitative analysis of CD3, CD8, CD45RO, Mac 387 Cytokine mRNA expression: IL-1 β , TNF, IL-2, IL-4, IL-6, IL-8, IL-10, GM-CSF, IFN γ Weighted kappa coefficient
	Musters <i>et al</i> 2018 ¹⁸	Comparing intra-articular variability of T-cell repertoire	13 RA patients	Arthroscopic biopsies (n=7) Knee; 2 anatomically different sites: SP and IP recesses	Number of TCR β clones Number of HEC Impact of the HECs on the total repertoire Overlap between the 25 most expanded TCR β clones (%) Chao-modified Sørensen index
	Kennedy <i>et al</i> 1988 ²³	Minimal number of tissue fragments required to limit sampling error in large joints	11 RA patients	Surgical biopsies (n=11)	H&E, absolute value of cell density per high power field. Number of tissue fragments to analyse to reduce sampling error
	Boyle <i>et al</i> 2003 ²⁴	Minimal number of tissue fragments required to limit sampling error in large joints	5 RA patients	Surgical biopsies (n=5) Hip or knee	PBMC equivalent (cellular equivalents of expression (CE)) number for synovial samples Relative expression (RE) of mRNA IL-1 β , IL-6, TNF- α , MMP-1 CE and GAPDH CE Number of biopsies required to limit sample error to <25% for each IL-1 β , IL-6, TNF- α , MMP-1
	Humby <i>et al</i> 2015 ¹¹	Minimal number of tissue fragments required to limit sampling error in small joints	RA patients ▲ Assessment cohort (n=35) ▲ Validation cohort (n=6)	Synovial biopsies (n=41) ▲ Assessment cohort: US-NB (n=35) ▲ Validation cohort: arthroscopic biopsies (n=3); surgical biopsies (n=3)	Semi quantitative analysis of CD20+, CD3 + and CD68+ cells in LL and SL % mean difference for different samples

Continued

Table 1 Continued

RQ	Author and year	Objective	Patients	Procedures	Outcomes
	Scire <i>et al</i> 2007 ²⁵	Minimal number of biopsies required to limit sampling error in small joints	RA patients ► Assessment cohort (n=9) ► Validation cohort (n=9)	Synovial biopsies (n=18) ► Assessment cohort: surgical biopsies (n=9) ► Validation cohort: US-NB (n=9)	Quantitative analysis of CD20+, CD3 + and CD68+ cells in LL and SL % mean difference for different samples or cutting levels
RQ5	Baeten <i>et al</i> 2000 ²⁶	Comparing in early vs established RA: ► Cell infiltrate	30 RA patients ► Early RA <1 year: (n=16) ► Established RA >1 year (n=14)	Arthroscopic biopsies (n=30)	H&E maximal LL thickness, mean LL thickness, vascularity, cell infiltrate Semi quantitative Integrines alphaVbeta3 and 5
	Tak <i>et al</i> 1997 ²⁷	Comparing in early vs established RA: ► Cell infiltrate	67 RA patients ► Early RA <1 year (n=31) ► Established RA >5 years (n=36)	Blind NB (n=67)	H&E LL thickness, vascularity, cell infiltrate immunohistochemistry (IHC): ► Semi quantitative analysis of CD68+ (LL and SL), CD3+, CD4+, CD8+, CD22+, CD38 +cells, mast cells ► Semi quantitative analysis of TNF α , IL-1 β and IL-6 +cells
	Katrib <i>et al</i> 2001 ⁴⁰	Comparing in early vs established RA: ► Cytokines expression ► MMPs expression	44 RA patients ► Early RA <1 year (n=22) ► Established RA >5 years (n=22)	Blind NB (n=44)	Quantitative CD68 +cells IHC: quantitative analysis of MCP-1, MIP-1alpha, MMP-1, MMP-3, TIMP-1, TIMP-2 by IHC
	Ospelt <i>et al</i> 2008 ²⁹ REF34	Comparing in early vs established RA: ► TLR expression	21 RA patients ► Early RA <1 year (n=10) ► Established RA >8 years (n=11)	Synovial biopsies n=44 ► Early RA: blind NB (n=10) ► Established RA: surgical biopsies (n=11)	IGC: semi quantitative expression of TLR2, 3 and 4
	Dubikov and Kalinichenko 2010 ³⁰	Comparing in early vs established RA: ► Apoptosis	78 RA patients ► Early RA <3 years (n=35) ► Established RA >3 years (n=43)	Arthroscopic and surgical biopsies (n=78)	Apoptosis (LL and SL) Quantitative DNA fragmentation staining positive cells p53 +cells and Bcl2 +cells
	Tak <i>et al</i> 1999 ²⁸	Comparing in early vs established RA: ► p53 expression	27 RA patients ► Early RA <6 months (n=13) ► Established RA >5 years (n=14)	Blind NB (n=27)	Semi quantitative evaluation of % of p53 +cells Western blot p53

Continued

Table 1 Continued

RQ	Author and year	Objective	Patients	Procedures	Outcomes
	Alivernini <i>et al</i> 2017 ³¹	Comparing in early vs remission and LDA in RA: ▲ Cell infiltrate	85 RA patients ▲ Remission (n=25) ▲ Low disease activity (n=10) ▲ Active Naïve (n=50)	US-NB (n=85)	Semiquantitative CD68+, CD3+, CD20 + and CD21+ cell infiltrate
	Rivellese <i>et al</i> 2020 ³³	Comparing in early naïve RA and established TNF inhibitors resistant RA: ▲ CD20 +infiltrate	329 RA patients ▲ Early RA; naïve (n=165) ▲ Established RA; TNF-inhibitors resistant (n=164)	Synovial biopsies (n=329) ▲ US-NB (n=325) ▲ Arthroscopic biopsies (n=14)	Semiquantitative scoring method (scale of 0–4) and digital image analysis (CD20 +area fraction) histologic B-cell scores
	Alivernini <i>et al</i> 2021 ³²	Comparing in early vs remission and low disease activity vs established RA: ▲ Krenn Synovitis Score	545 RA patients ▲ Early RA (n=240) ▲ Established RA (n=213) ▲ Remission RA (n=92)	US-NB (n=545)	Krenn Synovitis Score (lining layer thickness, stromal activation and cell infiltrate)
	Ostojic <i>et al</i> 2017 ³⁴	Comparing in early vs advanced stage osteoarthritis (OA): ▲ Grade of synovitis ▲ Expression of NF-kB and iNOS	20 OA patients ▲ Early radiographical OA (K-L 1–2) (n=10) ▲ Advanced radiographical OA (K-L 3–4) (n=10)	Synovial biopsies (n=20) ▲ Early radiographical: arthroscopic biopsies (n=10) ▲ Advanced radiographical: surgical biopsies (n=10)	Krenn Synovitis Score (lining layer thickness, stromal activation and cell infiltrate) Quantitative iNOS and NF-kB +cells per mm ²
	Guo <i>et al</i> 2018 ³⁶	Examine PD-1 pathway at various stages of RA development	152 RA patients ▲ Early RA <1 year (n=57) ▲ Established RA >1 year (n=95)	Synovial biopsies (n=152) ▲ Early RA: US-NB (n=57) ▲ Established RA: arthroscopic biopsies (n=95)	<i>PDCD1</i> , <i>CD274</i> and <i>PDCD1LG2</i> expression PD-1-PD-L1 pathway activity
	Guo <i>et al</i> 2017 ³⁶	Examine CD-40-CD40-L pathway in various stages of RA progression	152 RA patients Early RA <1 year (n=57) Established RA >1 year (n=95)	Synovial biopsies (n=152) Early RA: US-NB (n=57) Established RA: arthroscopic biopsies (n=95)	<i>CD40LG</i> and <i>CD40</i> expression CD40-CD40L pathway activity
	Lequerré <i>et al</i> 2009 ³⁸	Comparing in early vs established RA: ▲ Gene expression profile	8 RA patients ▲ Early RA <9 months (n=4) ▲ Established RA >4 years (n=4)	Arthroscopic biopsies (n=15)	Inflammatory pathways gene clusters and molecular signatures

Continued

Table 1 Continued

RQ	Author and year	Objective	Patients	Procedures	Outcomes
	Alivernini <i>et al</i> 2020 ³⁷	Compare in early vs established RA vs remission: ▲ Gene expression profile	112 RA patients ▲ Early naïve (n=45) ▲ Established resistant (n=31) ▲ Remission (n=36)	US-NB (n=112)	ST macrophage clusters tracked in ST using single-cell RNAseq across disease activity and stage

Bcl2, B-cell lymphoma 2; CD, cluster of differentiation; CPJ, cartilage-pannus junction; GAPDH, glyceraldehyde 3-phosphate dehydrogenase; GM-CSF, granulocyte-macrophage colony stimulating factor; IHC, immunohistochemistry; IL, interleukin; iNOS, inducible nitric oxide synthase; IP, infra-patellar recesses; JIA, juvenile idiopathic arthritis; LL, lining layer; MCP, metacarpophalangeal joint; MMP, matrix metallo-proteinases; NB, needle biopsy; OA, osteoarthritis; PBMC, peripheral blood mononuclear cell; P&F, portal and forceps; PsA, psoriatic arthritis; RA, rheumatoid arthritis; RIN, RNA integrity number; RQ, research question; SB, synovial biopsy; SL, sublining; SP, supra-patellar recess; SpA, spondyloarthritis; ST, synovial tissue; TCR, T-cell repertoire; TNF, tumour necrosis factor; UA, undifferentiated arthritis; US, ultrasound; VAS, visual analogic scale.

could not identify any difference in immune cell infiltrate and IL-6 expression assessed by non-parametric paired analysis.¹⁷ In addition, correlations were found for CD68 +macrophages, CD3 +T cells, CD138 +plasma cells and IL-6 expression and CD68 +macrophages and CD3 +T cells in the ST sublining of both small and large joints.¹⁷ As far as the synovial lining layer is concerned, significant correlations were found between small and large joints for lining layer hyperplasia, inflammatory infiltrate in H&E staining.¹⁹ The same transcripts were shown to be overexpressed in both small and large joints, and the same T-cell clone repartition was observed in small and large and right and left joints of the same individuals with RA.^{18 19}

It is noteworthy that all studies included small numbers of patients and, therefore, the absence of difference could also reflect a lack of power rather than a true similarity.

Impact of tissue handling and sample randomisation on tissue outcomes

One article with unclear RoB²⁰ assessed the impact of ST handling on quality and outcomes. They compared different tissue preservation (fresh vs cryopreserved tissue) and dissociation methods (mechanical vs mechanical +enzymatic) in tissue obtained from n=16 US-guided NB and P&F performed in RA patients and/or surgical biopsies performed n=32 RA patients. Interestingly, the cryopreservation did not impact the viable cell count or repartition of synovial cell subpopulations assessed by flow cytometry compared with freshly processed tissue. On the other hand, the mechanical +enzymatic technique using Liberase retrieved a higher total cell count per gram of ST, higher viable cell count and better repartition of cell subpopulations.

Few studies specifically assessed intra-articular variability. Authors retrieved ST from different compartments of the knee joint using arthroscopy, including comparisons of cartilage pannus-junction (CPJ) and supra-patellar recess (SP),²¹ lateral, medial and SP recesses versus CPJ²² and SP versus infra-patellar (IP) recess.¹⁸ The immune cell infiltrate assessed by IHC showed no major difference among the different compartments of the knee, including CD68+, CD3+, CD4+, CD8+, CD22+, CD38 + cells²¹ and CD45RO+, CD3+, CD8 + and Mac 387+ cells.²² These findings need to be interpreted carefully due to a low number of samples leading to a possible lack of power. Granzyme B and matrix metallo-proteinases (MMP-1, MMP-3 and MMP-13) expression was also assessed by IHC showing no differences.²¹ In addition, IL-1 β , TNF, IL-2, IL-4, IL-6, IL-8 and IFN γ levels were correlated in all knee recesses, while IL-10, granulocyte-macrophage colony stimulating factor (GM-CSF) did not appear correlated across knee compartments.²² The TCR repertoire study showed no difference in the most frequently expressed T-lymphocyte clones across SP and IP compartments.¹⁸

Table 2 Risk of bias assessment of all included studies

RQ	Study	Study desing	Q1 inclusion criteria	Q2 study population	Q3 exposure	Q4 objective	Q5 confounding factors	Q6 confounding factors	Q7 outcomes measure	Q8 statistical analysis	Number of ✓	RoB
RQ1	Humby <i>et al</i> ¹⁵	Retrospective cohort	✓	✓	✓	✓	×	×	✓	✓	6/8	Unclear
	Just <i>et al</i> ¹⁶	Retrospective cohort	✓	✓	✓	✓	✓	×	✓	✓	7/8	Unclear
RQ2	Kraan <i>et al</i> ¹⁷	Cross-sectional	✓	×	✓	✓	×	×	✓	✓	6	High
	Musters <i>et al</i> ¹⁸	Cross-sectional	✓	✓	✓	×	×	×	✓	✓	5	High
RQ3	Triaille <i>et al</i> ¹⁹	Cross-sectional	✓	✓	✓	✓	×	×	✓	✓	6	Unclear
	Donlin <i>et al</i> ²⁰	Cross-sectional	✓	×	✓	✓	×	×	✓	✓	5	High
	Smeets <i>et al</i> ²¹	Cross-sectional	✓	✓	✓	✓	×	×	✓	✓	6	Unclear
	Kirkham <i>et al</i> ²²	Cross-sectional	✓	✓	✓	✓	×	×	✓	✓	6	Unclear
	Kennedy <i>et al</i> ²³	Cross-sectional	✓	×	✓	✓	×	×	✓	✓	5	High
	Boyle <i>et al</i> ²⁴	Cross-sectional	✓	×	✓	✓	×	×	✓	✓	5	High
	Humby <i>et al</i> ¹¹	Retrospective cohort	✓	✓	✓	✓	×	×	✓	✓	6	Unclear
	Musters <i>et al</i> ¹⁸	Cross-sectional	✓	✓	✓	×	×	×	✓	✓	5	High
RQ5	Scire <i>et al</i> ²⁵	Cross-sectional	✓	✓	✓	✓	×	×	✓	✓	6	Unclear
	Beaten <i>et al</i> ²⁶	Cross-sectional	✓	×	✓	✓	×	×	✓	✓	5	High
	Tak <i>et al</i> ²⁷	Cross-sectional	✓	✓	✓	✓	×	×	✓	✓	6	Unclear
	Tak <i>et al</i> ²⁸	Cross-sectional	✓	×	✓	✓	×	×	✓	✓	5	High
	Ospelt <i>et al</i> ²⁹	Cross-sectional	✓	×	✓	✓	×	×	✓	✓	5	High
	Dubikov and Kalinichenko ³⁰	Cross-sectional	✓	×	✓	✓	×	×	✓	✓	5	High
	Alivernini <i>et al</i> ³¹	Cross-sectional	✓	✓	✓	✓	✓	×	✓	✓	7	Unclear
	Alivernini <i>et al</i> ³²	Cross-sectional	✓	✓	✓	✓	✓	×	✓	✓	7	Unclear
	Rivellese <i>et al</i> ³³	Cross-sectional	✓	✓	✓	✓	✓	×	✓	✓	7	Unclear
	Katrib <i>et al</i> ⁴⁰	Cross-sectional	✓	×	✓	✓	×	×	✓	✓	5	High
	Ostojic <i>et al</i> ³⁵	Cross-sectional	✓	✓	✓	✓	✓	×	✓	✓	7	Unclear
	Guo <i>et al</i> ³⁵	Cross-sectional	✓	×	✓	✓	×	×	✓	✓	5	High
	Guo <i>et al</i> ³⁶	Cross-sectional	✓	×	✓	✓	×	×	✓	✓	5	High
	Lequerré <i>et al</i> ³⁷	Cross-sectional	✓	✓	✓	✓	✓	×	✓	✓	7	Unclear
	Alivernini <i>et al</i> ³⁸	Cross-sectional	✓	✓	✓	✓	✓	×	✓	✓	7	Unclear

RoB, risk of bias; RQ, research question; Q, question

Table 3 Reporting of different areas pertaining to synovial tissue sampling, handling or analysing

		N	%
Clinical data	Age and RMDs	26/26	100
	Disease duration	24/26	92
	Disease activity	16/26	62
	Treatments	16/26	62
	US features (if US-guided SB)	4/11	36
Biopsy procedure	Target joint(s)	21/26	81
	Operators' experience	0/24	0
	Adverse events	5/24	20
Tissue handling	Number of fragments	19/26	73
	Processing: fixation, freezing, fresh use, specific processing details	26/26	100
Tissue quality control		17/26	65
Tissue outcomes	Imaging: scoring/quantification, area assessed	25/26	96
	Reference to another publication for scoring system	8/26	30
	Intraobserver/interobserver reproducibility	4/26	15
	RNA/DNA: purity, quality, analysis	4/10	40
	Disaggregation: isolation method, purity and viability	2/2	100

Green, 71–100%; Orange, 36–70%; Red, 0–35%; RMDs, rheumatic and musculoskeletal diseases; SB, synovial biopsy; US, ultrasound.

Finally, a few studies with high^{23 24} and unclear RoB^{15 25} studied sampling error by evaluating the minimum number of tissue samples needing to be included in a single analysis to obtain consistent results. Of note, two of these studies assessed ST retrieved from arthroplasty surgery and not synovial biopsies. In small joints, it was demonstrated that the semiquantitative IHC analysis of four tissue fragments provided a reliable sample analysis with 10% sampling error (mean difference <10% in 91.6% of the cases).¹¹ If quantitative analysis is used, the analysis of a cumulative area of 2.5 mm² and/or six (large joints) to eight (small joints) randomly selected sections (from different samples or from different cutting levels) allowed a mean difference <10% in 100% of cases.^{23 25} As far as cytokine expression (IL-1 β , IL-6, TNF- α and MMP-1) by PCR in large joints is concerned, four to seven tissue fragments

were deemed necessary in order to detect a twofold change with a 25% sampling error.²⁴

Impact of the disease stage on the tissue outcomes

Fourteen studies assessed the impact of disease stage on histopathological findings in RA,^{26–33} osteoarthritis (OA)³⁴ or transcriptomics aspects in RA,^{35–38} including eight studies with high RoB and seven with unclear RoB.^{2 27 31–34 37} Of interest, while contradictory results for lining layer thickness and CD68 + cell infiltrate (lining layer and sublining) in early versus established RA were reported, other cell infiltrates (CD3, CD4, CD8, CD20 or CD22, CD38 and mast cells) and vascularity in sublining appeared to be the same between early and established RA.^{26 39} In addition, proinflammatory cytokines (TNF α , IL-1 β , IL-6 and MCP-1), MMPs, toll-like receptors 2, 3 and 4 and integrins α V β 5 and 3 did not appear to be expressed differentially in STs across disease stages,^{26 29 39 40} while cells displaying apoptotic features and p53 positive cells were increased in established RA.^{28 30} Another study has compared histopathological infiltrate between knee synovitis in patient with active naïve RA and US power Doppler negative knee synovitis in patients with RA in remission or low disease activity. Interestingly, authors showed a significantly higher infiltrate of CD68+, CD20+, CD21 + and CD3+ cell in ST from naïve RA patients.³¹ Two other studies interrogated ST infiltrate in early naïve versus resistant established RA versus RA in remission.^{32 33} In one study, the authors report that the Krenn Synovitis Score (KSS) was higher in patients with early active than established RA and both were higher compared with patients in remission.³² In addition, established RA resistant to TNFi had higher KSS and higher prevalence of B cell-rich synovitis than early untreated RA.³³ It is important to note that the definition of 'early' and 'established' RA was variable across studies and this might explain the contradictory results observed. Detailed results are summarised in tables 6 and 7.

As far as OA is concerned, Ostojic *et al* have described an increased lining layer thickness and stromal infiltration in mild OA (Kellgren and Lawrence Score (KLS) 1–2) compared with advanced OA (KLS 3–4), while cell infiltrate was more pronounced in advanced OA compared with mild. Overall, the KSS was superior in mild OA suggesting synovitis in early stages compared with more advanced and severe OA (table 7).³⁴

Finally, four studies looked into transcriptomic expression across the RA spectrum. The two studies from Guo *et al* showed no difference in expression of genes related to CD40–CD40L³⁶ or PD-1 PD-L1³⁵ pathways activity in STs from 57 early RA (<1 year) versus 95 established RA (>1 year). Lequerré *et al* showed differing transcriptomic signatures in a lower number of samples (n=3 early RA (<9 months) and n=4 established RA (>4 years)).³⁷ Early RA samples overexpressed 503 genes corresponding to

Table 4 Tissue outcome and tolerance of synovial biopsies

	US-NB*	US-P&F*	NB*
Tissue yield and quality ¹⁵	=	=	Number of graded ST fragments/ total number of ST fragments <Small joints trend p=0.057 <Large joint p=0.048
Large joints RNA yield (quantity) ¹⁵	A>US-NB: p=0.002	A>US-P&F: p=0.00REF	–
Large joints RNA quality (RIN) ¹⁵	A>US-NB: p=0.0018	A=US-P&F: p=0.068	–
Small and large joints Immune cell infiltrate and pathotype ¹⁵	=	=	=
Tolerance VAS pain, swelling, stiffness ¹⁶	=	=	–
Willingness to repeat ¹⁶	US-P&F>US-NB: p<0.01 A>US-NB: p<0.01	=	–

*Arthroscopy is the reference in this table.

=, no statistical difference; –, not done; A, arthroscopy; NB, needle biopsy; P&F, portal and forceps; RIN, RNA integrity number; ST, synovial tissue; US, ultrasound; VAS, visual analogic scale.

12 biological processes, including immunity and host defenses, stress responses, T-cell-mediated immunity and tumour suppressor and MHC class II mediated immunity; while established RA tissues overexpressed 216 genes linked to the following biological processes: cell cycle, cell surface receptor-mediated signal transduction, cell cycle control, ligand-mediated signalling, apoptosis inhibition and granulocyte-mediated immunity. In addition, Alivernini *et al* assessed distinct ST macrophage clusters using single-cell RNA sequencing in tissues collected by US-guided biopsies across different RA phases (naïve RA vs established RA vs remission RA, respectively). Of interest, MerTK +clusters (TREM2high and LYVE1pos)

and increased proportions of MerTK-CD48-SPP1+ and MerTK-CD48-S100A12+ clusters were observed in naïve and resistant RA, while ST from patients with RA in remission were characterised by an increase in the MerTKposFOLR2highLYVE1pos cluster. Overall, these data are suggestive of different ST macrophages clusters with very diverse phenotypes and function acting at different periods of the disease course.³⁸

DISCUSSION

This SLR summarises the available evidence on the impact of several aspects of SB methods on tissue

Table 5 Tissue outcome and tolerance of sequential synovial biopsies

	US-NB*	US-P&F*	A*
Tissue yield and quality: ► Proportion of graded ST fragments ► Total graded ST area ► ST area per graded biopsy fragment ¹⁵	=	=	=
RNA yield Small joints	US-NB1 >US-NB2: p=0.026	–	–
RNA yield Large joints ¹⁵	–	US-P&F1>USP&F2: p<0.001	A1 <A2: p<0.001
RNA quality (RIN) Small and large joints ¹⁵	=	=	=
Immune cell infiltrate and pathotype ¹⁵	=	=	=
Tolerance VAS pain, swelling, stiffness ¹⁶	=	=	=
Willingness to repeat ¹⁶	=		
Adverse events ¹⁶	=		

*First biopsy is the reference in this table.

A, arthroscopy; NB, needle biopsy; P&F, portal and forceps; RIN, RNA integrity number; ST, synovial tissue; US, ultrasound; VAS, visual analogic scale

Table 6 Impact of disease stage on histopathological tissue outcomes in RA

Early vs established RA			
	Early=established	Early >established	Established >early RA
H&E	Mean lining layer thickness ^{26 27} Vascularity ²⁶	Mean LL thickness ^{26 27}	Maximum LL thickness ²⁶
Cell infiltrate	Mast cells ²⁷ Semiquantitative CD68 +cells ²⁷ CD3, CD4, CD8, CD20 or CD22 and CD38 +cells ²⁷	Bcl2 +cells ³⁰	CD68 +cells ³⁴ p53 +cells ³⁰
Cytokines	TNFα, IL- 1β, IL-6 ²⁷ MCP-1, MMP-1 and 3, TIMP-1 and 2 ³⁴	–	MIP-1α ³⁴
Proteins	TCR 2, 3, 4 ²⁹ p53 protein expression (tissue lysates) ²⁷ Integrines αVβ5 and 3 ²⁶	–	–

Bcl2, B-cell lymphoma 2; CD, cluster of differentiation; GM-CSF, granulocyte-macrophage colony stimulating factor; IFN, interferon; IHC, immunohistochemistry; IL, interleukin; LL, lining layer; MCP, monocyte chemoattractant protein-1; MI-1α, macrophage inflammation protein 1α; MMP, matrix metallo-proteinase; RA, rheumatoid arthritis; TCR, T-cell repertoire; TIMP, tissue inhibitor of metalloproteinases; TNF, tumour necrosis factor.

outcomes. To the best of our knowledge, this is the first systematic approach examining the impact of a wide range of methods on the results of ST assessment, whether it is histological, bulk transcriptomic or more advanced single cell technologies. We were able to retrieve a number of studies suggesting that the biopsy technique, regardless of the guiding technique (arthroscopic or US guided) was well tolerated and allowed to retrieve a sufficient amount of tissue for a meaningful analysis, suggesting that any of these techniques can be used for translational research. On the other hand, tissue yield and quality were consistently shown to be reduced with blind NB technique. The analysis also showed that most histopathological and transcriptomic outcomes were similar regardless of the joint biopsied (small vs large, right vs left) or the recess of the joint that was biopsied in the same patient (ie, knee recesses). Of note, all the studies pertaining to this RQ were performed in ST retrieved in RA patients and these results are not necessarily applicable to other RMDs. Similar studies using the most advanced technologies are however warranted, since joint-dependent epigenetic differences among ST resident cells have been reported.⁴¹

It is noteworthy that a low number of studies was retrieved to address the different RQs, and one RQ pertaining to the impact of tissue quality on outcomes was not addressed due to the absence of relevant articles. Moreover, it is important to mention that the majority of the studies was performed in patients with

RA and in research settings; and those findings may not be necessarily generalisable to other inflammatory conditions since the pathogenetic mechanisms may be different, or other clinical settings. In addition, all the articles had a high or unclear RoB, frequently due to the existence of confounding factors. These biases related mostly to the study design that may affect the results. Therefore, efforts should be undertaken to reduce them. This could be achieved by carefully designing studies with more homogeneous inclusion criteria and better-defined baseline populations. While in the past the paucity of centres with SB expertise and the invasiveness of the procedures were a limiting factor, the implementation of US-guided SB procedures in a larger number of centres, thanks to an improved tolerance and reduced invasiveness, will help in evaluating the real impact of this procedure. In addition, it will allow the development of prospective studies with a higher number of patients better powered to take into account both clinical and tissue heterogeneity.

Although this SLR aimed to assess evidence in both clinical and research settings, the studies referring to clinical practice were extremely scarce. Therefore, the analysis of ST should be reinforced as an area of high interest and studies aimed at defining disease endotypes and their applications, both for research purposes, and also for practicing clinicians, should be prioritised over the coming years. Such studies will contribute to building data sets and evidence

Table 7 Impact of disease stage on histopathological tissue outcomes in OA

Early vs advanced OA			
	Early=advanced	Early >advanced OA	Advanced>early OA
Cell infiltrate	–	Krenn Synovitis Score ³⁵	–
Others	iNOS (IHC) (sublining) ³⁵	NF-κB (IHC) ³⁵	–

IHC, immunohistochemistry; iNOS, inducible nitric oxide synthase; NF-κB, nuclear factor kappa-light-chain-enhancer of activated B cells; OA, osteoarthritis.

supporting the use of ST analysis for standard of care diagnosis, prognosis or disease management.

In conclusion, this SLR retrieved evidence which contributed to inform the EULAR PtC for minimal reporting requirements in ST research in rheumatology.

Author affiliations

¹Musculoskeletal Research Group, Institute of Infection, Immunity and Inflammation, University of Glasgow College of Medical Veterinary and Life Sciences, Glasgow, UK

²Université Paris-Saclay, UVSQ, Inserm U1173, Infection et inflammation, Laboratory of excellence Inflamex, Paris, Île-de-France, France

³Rheumatology Department, Ambroise Paré Hospital, Boulogne-Billancourt, France

⁴Medicine, University of Paris Health Interuniversity Library, Paris, Île-de-France, France

⁵Rheumatology Research Group and Research into Inflammatory Arthritis Centre Versus Arthritis, Institute of Inflammation and Ageing, University of Birmingham, Birmingham, UK

⁶Rheumatology, Fondazione Policlinico Universitario Agostino Gemelli IRCCS, Roma, Lazio, Italy

Twitter Aurélie Najm @AurelieRheumo

Contributors All authors contributed to and finally approved the current manuscript. MAD'A and AF share last authorship. AN acts as guarantor.

Funding This work was funded by EULAR (SCI020).

Competing interests AN has received consulting and/or speaker's fees from UCB, CHUGAI, BMS all unrelated to this manuscript. FC has received consulting and/or speaker's fees from Lilly, Novartis and UCB, all unrelated to this manuscript. MAD'A has received consulting and/or speaker's fees from Lilly, Novartis, AbbVie, BMS, Pfizer, Galapagos and UCB, all unrelated to this manuscript. AF has received consulting, research grants and/or speaker's fees from Abbvie, Roche, Janssen, UCB, Nascient, Mestag, GSK, all unrelated to this manuscript.

Patient consent for publication Not required.

Ethics approval This study does not involve human participants.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement All data relevant to the study are included in the article or uploaded as supplementary information. Not applicable.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Aurélie Najm <http://orcid.org/0000-0002-6008-503X>

Félicie Costantino <http://orcid.org/0000-0002-1449-959X>

Andrew Filer <http://orcid.org/0000-0001-8164-6249>

REFERENCES

- Croft AP, Campos J, Jansen K, *et al*. Distinct fibroblast subsets drive inflammation and damage in arthritis. *Nature* 2019;570:246–51.
- Alivernini S, MacDonald L, Elmesari A. Distinct synovial tissue macrophage subsets regulate inflammation and remission in rheumatoid arthritis. *Nat Med* 2021.
- Orange DE, Donlin LT. Advancing rheumatoid arthritis synovial biopsy analysis: one B cell at a time. *Arthritis Rheumatol* 2020;72:702–3.
- Wade SM, Canavan M, McGarry T, *et al*. Association of synovial tissue polyfunctional T-cells with DAPSA in psoriatic arthritis. *Ann Rheum Dis* 2019;78:350–4.
- Zhang F, Wei K, Slowikowski K, *et al*. Defining inflammatory cell states in rheumatoid arthritis joint synovial tissues by integrating single-cell transcriptomics and mass cytometry. *Nat Immunol* 2019;20:928–42.
- Najm A, Orr C, Heymann M-F, *et al*. Success rate and utility of ultrasound-guided synovial biopsies in clinical practice. *J Rheumatol* 2016;43:2113–9.
- Dyke B, Maybury M, Alivernini S. Survey of Minimally-Invasive Ultrasound-Guided Synovial Biopsy Performed at Rheumatology Centres [abstract]. *Arthritis Rheumatol* 2020;72 <https://acrabstracts.org/abstract/survey-of-minimally-invasive-ultrasound-guided-synovial-biopsy-performed-at-rheumatology-centres/>
- Romao VC, Polido-Pereira J, Barros R. Ultrasound-guided synovial needle biopsy: single center experience of an emerging, minimally invasive technique in clinical practice and research. *Ann Rheum Dis* 2017;76:746–7.
- Kelly S, Bombardieri M, Humby F. Ultrasound guided synovial biopsies: safety, tolerability, and tissue quality. *Rheumatology* 2012;51:ii68.
- Lazarou I, Kelly SG, Meric de Bellefon L. Ultrasound-Guided synovial biopsies of wrists, metacarpophalangeal, Metatarsophalangeal, interphalangeal joints, and tendon sheaths. *Front Med* 2019;6:2.
- Humby F, Kelly S, Hands R, *et al*. Use of ultrasound-guided small joint biopsy to evaluate the histopathologic response to rheumatoid arthritis therapy: recommendations for application to clinical trials. *Arthritis Rheumatol* 2015;67:2601–10.
- van der Heijde D, Aletaha D, Carmona L, *et al*. 2014 update of the EULAR standardised operating procedures for EULAR-endorsed recommendations. *Ann Rheum Dis* 2015;74:8–13.
- Eriksen MB, Frandsen TF. The impact of patient, intervention, comparison, outcome (PICO) as a search strategy tool on literature search quality: a systematic review. *J Med Libr Assoc* 2018;106:420–31.
- Martin J. Joanna Briggs Institute 2017 critical appraisal checklist for analytical cross sectional studies. 7, 2017.
- Humby F, Romão V, Manzo A. A multi centre retrospective analysis evaluating performance of synovial biopsy techniques: arthroscopic vs. ultrasound guided vs. blind needle biopsy. *Arthritis Rheumatol* 2018;70.
- Just SA, Humby F, Lindegaard H, *et al*. Patient-Reported outcomes and safety in patients undergoing synovial biopsy: comparison of ultrasound-guided needle biopsy, ultrasound-guided portal and forceps and arthroscopic-guided synovial biopsy techniques in five centres across Europe. *RMD Open* 2018;4:e000799.
- Kraan MC, Reece RJ, Smeets TJM, *et al*. Comparison of synovial tissues from the knee joints and the small joints of rheumatoid arthritis patients: implications for pathogenesis and evaluation of treatment. *Arthritis Rheum* 2002;46:2034–8.
- Musters A, Klarenbeek PL, Doorenspleet ME, *et al*. In rheumatoid arthritis, synovitis at different inflammatory sites is dominated by shared but patient-specific T cell clones. *J Immunol* 2018;201:1800421:417–22.
- Triaille C, Vansteenkiste L, Constant M, *et al*. Paired Rheumatoid Arthritis Synovial Biopsies From Small and Large Joints Show Similar Global Transcriptomic Patterns With Enrichment of Private Specificity TCRB and TCR Signaling Pathways. *Front Immunol* 2020;11:593083.
- Donlin LT, Rao DA, Wei K, *et al*. Methods for high-dimensional analysis of cells dissociated from cryopreserved synovial tissue. *Arthritis Res Ther* 2018;20:139.
- Smeets TJM, Barg EC, Kraan MC, *et al*. Analysis of the cell infiltrate and expression of proinflammatory cytokines and matrix metalloproteinases in arthroscopic synovial biopsies: comparison with synovial samples from patients with end stage, destructive rheumatoid arthritis. *Ann Rheum Dis* 2003;62:635–8.
- Kirkham B, Portek I, Lee CS, *et al*. Intraarticular variability of synovial membrane histology, immunohistology, and cytokine mRNA expression in patients with rheumatoid arthritis. *J Rheumatol* 1999;26:777–84.
- Kennedy TD, Plater-Zyberk C, Partridge TA, *et al*. Representative sample of rheumatoid synovium: a morphometric study. *J Clin Pathol* 1988;41:841–6.
- Boyle DL, Rosengren S, Bugbee W, *et al*. Quantitative biomarker analysis of synovial gene expression by real-time PCR. *Arthritis Res Ther* 2003;5:R352–60.
- Scirè CA, Epis O, Codullo V, *et al*. Immunohistological assessment of the synovial tissue in small joints in rheumatoid arthritis: validation of a minimally invasive ultrasound-guided synovial biopsy procedure. *Arthritis Res Ther* 2007;9:R101.
- Baeten D, Demetter P, Cuvelier C, *et al*. Comparative study of the synovial histology in rheumatoid arthritis, spondyloarthritis, and osteoarthritis: influence of disease duration and activity. *Ann Rheum Dis* 2000;59:945–53.
- Tak PP, Smeets TJM, Daha MR, *et al*. Analysis of the synovial cell infiltrate in early rheumatoid synovial tissue in relation to local disease activity. *Arthritis & Rheumatism* 1997;40:217–25.
- Tak PP, Smeets TJ, Boyle DL, *et al*. P53 overexpression in synovial tissue from patients with early and longstanding rheumatoid arthritis

- compared with patients with reactive arthritis and osteoarthritis. *Arthritis Rheum* 1999;42:948–53.
- 29 Ospelt C, Brentano F, Rengel Y, *et al.* Overexpression of Toll-like receptors 3 and 4 in synovial tissue from patients with early rheumatoid arthritis: Toll-like receptor expression in early and longstanding arthritis. *Arthritis Rheum* 2008;58:3684–92.
 - 30 Dubikov AI, Kalinichenko SG. Small molecules regulating apoptosis in the synovium in rheumatoid arthritis. *Scand J Rheumatol* 2010;39:368–72.
 - 31 Alivernini S, Tolusso B, Petricca L, *et al.* Synovial features of patients with rheumatoid arthritis and psoriatic arthritis in clinical and ultrasound remission differ under anti-TNF therapy: a clue to interpret different chances of relapse after clinical remission? *Ann Rheum Dis* 2017;76:1228–36.
 - 32 Alivernini S, Tolusso B, Gessi M, *et al.* Inclusion of synovial tissue-derived characteristics in a nomogram for the prediction of treatment response in treatment-naïve rheumatoid arthritis patients. *Arthritis Rheumatol* 2021;73:1601–13.
 - 33 Rivellese F, Humby F, Bugatti S, *et al.* B cell synovitis and clinical phenotypes in rheumatoid arthritis: relationship to disease stages and drug exposure. *Arthritis Rheumatol* 2020;72:714–25.
 - 34 Ostojic M, Soljic V, Vukojevic K, *et al.* Immunohistochemical characterization of early and advanced knee osteoarthritis by NF- κ B and iNOS expression. *J Orthop Res* 2017;35:1990–7.
 - 35 Guo Y, Walsh AM, Canavan M, *et al.* Immune checkpoint inhibitor PD-1 pathway is down-regulated in synovium at various stages of rheumatoid arthritis disease progression. *PLoS One* 2018;13:e0192704.
 - 36 Guo Y, Walsh AM, Fearon U, *et al.* CD40L-Dependent pathway is active at various stages of rheumatoid arthritis disease progression. *J Immunol* 2017;198:4490–501.
 - 37 Lequerré T, Bansard C, Vittecoq O, *et al.* Early and long-standing rheumatoid arthritis: distinct molecular signatures identified by gene-expression profiling in synovia. *Arthritis Res Ther* 2009;11:R99.
 - 38 Alivernini S, MacDonald L, Elmesmari A, *et al.* Distinct synovial tissue macrophage subsets regulate inflammation and remission in rheumatoid arthritis. *Nat Med* 2020;26:1295–306.
 - 39 Tak PP, Smeets TJ, Daha MR, *et al.* Analysis of the synovial cell infiltrate in early rheumatoid synovial tissue in relation to local disease activity. *Arthritis Rheum* 1997;40:217–25.
 - 40 Katrib A, Tak PP, Bertouch JV, *et al.* Expression of chemokines and matrix metalloproteinases in early rheumatoid arthritis. *Rheumatology* 2001;40:988–94.
 - 41 Frank-Bertoncelj M, Trenkmann M, Klein K, *et al.* Epigenetically-driven anatomical diversity of synovial fibroblasts guides joint-specific fibroblast functions. *Nat Commun* 2017;8:14852.