LETTER

Characteristics of Sjögren's syndrome
associated with rheumatoid arthritisNolan Hassold ¹, Raphaele Seror ¹, Xavier Mariette ^{1,2},
Gaetane Nocturne^{1,3}

To cite: Hassold N, Seror R, Mariette X, *et al*. Characteristics of Sjögren's syndrome associated with rheumatoid arthritis. *RMD Open* 2022;**8**:e002234. doi:10.1136/rmdopen-2022-002234

Accepted 25 January 2022

© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

¹Department of Rheumatology, Hôpital Bicêtre, Le Kremlin-Bicêtre, France

²Université Paris-Sud, Center for Immunology of Viral Infections and Auto-immune Diseases (IMVA), Institut pour la Santé et la Recherche Médicale (INSERM) UMR 1184, Université Paris-Saclay, Le Kremlin Bicêtre, France

³Center for Immunology of Viral Infections and Autoimmune Diseases, Le Kremlin-Bicêtre, France

Correspondence to

Dr Nolan Hassold;
nolan.hassold-rugolino@aphp.fr

Sjögren Syndrome (SS) classically associates sicca syndrome, pain and asthenia. Systemic manifestations may develop with potentially severe organ damage including pulmonary, neurological, renal involvement and lymphoma.^{1,2} SS can be associated with rheumatoid arthritis (RA). Despite this frequent association, few studies have investigated differences between primary Sjögren syndrome (pSS) and SS associated with RA (SS/RA).^{3,4}

We have conducted a retrospective single-centre case-controls study. Patients were identified in the Paris-Saclay cohort, a prospectively collected database. Cases were patients fulfilling both ACR/EULAR 2016 criteria for SS and ACR 2010 criteria for RA and followed between 2000 and 2019. For each case, two controls from our local pSS cohort were randomly assigned and matched on disease duration. Disease activity was assessed using the EULAR Sjögren's Syndrome Disease Activity Index (ESSDAI)⁵ and ESSDAI cumulative score (Cumulative ESSDAI). The latter was calculated considering for each domain, the highest value reached by the patient during his/her follow-up. SS/RA and pSS patients were compared using a Chi-square test for non-normal binary variables and Mann-Whitney-Wilcoxon test for non-normal numeric variables. The p was retained as significant in the statistical analyses if <0.05.

Sixty-two SS/RA patients were compared with 124 pSS patients (table 1). There were more women among the pSS group (p=0.03). We observed a higher disease activity in the SS/RA group when considering ESSDAI and cumulative ESSDAI. We also performed analyses excluding joint involvement: there was no more difference between the two groups, which reflects the weight of the articular domain in the activity of the disease in the SS/RA group. There

was higher prevalence of arthralgia and arthritis in the SS/RA group. Pulmonary involvement was also significantly more frequent in SS/RA group; 9/62 (14.5%) patients (six interstitial lung disease (ILD): 4 non-specific interstitial pneumonia (NSIP) and two non-specific ILD), 1 bronchiectasis, 1 bronchial disorder and one pulmonary cysts) were compared with 7/123 (5.7%) patients in the pSS group (3 ILD: 2 NSIP and one non-specific ILD, 3 bronchiectasis and one obstructive bronchial disorder). pSS patients presented more skin or vaginal dryness and parotid gland enlargement. We found higher ESR levels and higher rate of RF positivity in the SS/RA group compared with pSS. Even though numerically higher in the pSS group, the frequency of lymphoma did not significantly differ between groups (p=0.267). There was no difference between the two groups on the other parameters.

Our study confirms previous reports showing that pSS patients have more frequently parotid gland enlargement.^{4,6} As described by He *et al*, we found that patients with SS/RA have more frequently articular and pulmonary involvement.⁴ It is difficult to decipher if articular involvement was linked to RA or to SS in these patients. Of note, in our study, more than 50% of the RA/SS patients had erosions. Regarding lung involvement, discrimination between SS linked or RA linked manifestations is also challenging. ILD complicating pSS tends to give more NSIP while ILD complicating RA tends to give usual interstitial pneumonia (UIP). In our study, pulmonary involvement seemed to be mainly due to SS in the SS/RA group since there was no UIP.

This study shows that patients with pSS and SS/RA have the same level of systemic disease activity. Thus, in RA patients, sicca symptoms should not be neglected and should be

Table 1 Comparison of pSS and SS/RA

	pSS	SS/RA	P value
Number of patients	124	62	
Female sex (n, %)	112/124 (90%)	49/62 (79%)	0.033
Age	55.06(13,9)	53.55(14,3)	0.539
Disease duration	8.8 years(8.8)	8.1 years (8.8)	
Tobacco	19/112 (17%)	16/55 (29%)	0.07
Asthenia (n, %)	99/122 (81%)	35/41 (85%)	0.541
Dry eye (n, %)	107/124 (86%)	47/56 (84%)	0.676
Dry mouth (n, %)	111/124 (89%)	48/55 (87%)	0.660
Others sicca syndrome (vagina and skin) (n, %)	90/118 (76%)	30/51 (58%)	0.022
Parotid enlargement (n, %)	61/123 (49.6%)	13/59 (22%)	<0001
Myositis (n, %)	1/123 (1%)	1/60 (2%)	0.602
Arthralgia (n, %)	61/108 (56.5%)	54/62 (87%)	<0001
Arthritis (n, %)	20/123 (16.3%)	57/62 (92%)	<0001
Peripheral neurological involvement (n, %)	10/121 (8.3%)	1/62 (1.6%)	0.073
Central neurological involvement (n, %)	2/123 (1.6%)	0/62 (0%)	0.313
Pulmonary involvement (n, %)	7/123 (5.7%)	9/62 (14.5%)	0.044
Renal involvement (n, %)	3/123 (2.4%)	0/61 (0%)	0.219
Lymphadenopathy (n, %)	15/123 (12.2%)	7/62 (11.3%)	0.858
Lymphoma (n, %)	9/123 (7%)	2/62 (3%)	0.267
IgG levels (g/L)(Mean (SD))	13.6 (6.6)/113	15 (4.7)/45	0.067
Presence of monoclonal peak (n, %)	10/114 (9%)	7/55 (13%)	0.423
CRP (mg/L)(Mean (SD))	6.3 (6.5)/112	13.6 (23.4)/61	0.184
ESR (mm at 1 hour)(Mean (SD))	23.6 (21)/109	39.7 (27.3)/58	<0001
Lymphocytes count (/mm ³)(Mean (SD))	1673 (1212)/118	1657(636)/62	0.494
Beta2-microglobulin (mg/L)(Mean (SD))	2.3 (1.1)/101	2.7 (1.5)/39	0.172
LDH (U/L)(Mean (SD))	344 (101)/105	350 (57.9)/37	0.344
C3 (g/L)(Mean (SD))	1.09 (0.3)/111	1.1 (0.3)/43	0.893
C4 (g/L)(Mean (SD))	0.31 (0.9)/111	0.22 (0.1)/42	0.987
Anti-SSA antibodies (n, %)	86/120 (72%)	35/60 (58%)	0.072
Anti-SSB antibodies (n, %)	48/122 (39%)	17/58 (29%)	0.19
RF positivity (n, %)	57/119 (48%)	54/62 (87%)	<0001
RF levels when detectable (UI/mL)(Mean (SD))	426(794)	348(569)	0.651
Anti-CCP (n, %)	1/107 (1%)	55/62 (89%)	<0001
Cryoglobulinemia (n, %)	2/123 (2%)	3/62 (5%)	0.203
ESSDAI	3.74 ⁵	6.3 (5.5)	<0001
Cumulative ESSDAI	6.8 (7.5)	9.2 (5.7)	<0001
ESSDAI without joint domain	2.85 (4.8)	3.2 (4.8)	0.971
Cumulative ESSDAI without joint domain	5.4 (7.4)	4.5 (5.3)	0.573

Variables are presented as mean (SD) or number and percentage.

explored in order to detect an associated SS and to adapt the follow-up accordingly especially concerning pulmonary involvement and risk of lymphoma.

Contributors All of the authors have contributed substantially to the manuscript, agree with the data and the interpretations of the manuscript content, and have given their permission to submit the attached version of the manuscript. None of the authors have conflicts of interest with any aspect of the manuscript. This manuscript has not been published elsewhere and, if accepted in your journal, will not be republished in any other journal in the same or similar form without consent from the Editor of RMD Open.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Consent obtained directly from patient(s)

Ethics approval This study involves human participants and was approved by local ethic committee Participants gave informed consent to participate in the study before taking part.

Provenance and peer review Not commissioned; externally peer reviewed.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which

permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Nolan Hassold <http://orcid.org/0000-0003-0944-4997>

Raphael Seror <http://orcid.org/0000-0002-5523-1856>

Xavier Mariette <http://orcid.org/0000-0002-4244-5417>

REFERENCES

- 1 Mariette X, Criswell LA. Primary Sjögren's syndrome. *N Engl J Med* 2018;378:931–9.
- 2 Nocturne G, Mariette X. Sjögren syndrome-associated lymphomas: an update on pathogenesis and management. *Br J Haematol* 2015;168:317–27.
- 3 Kollert F, Fisher BA. Equal rights in autoimmunity: is Sjögren's syndrome ever 'secondary'? *Rheumatology* 2020;59:1218–25.
- 4 He J, Ding Y, Feng M, *et al*. Characteristics of Sjögren's syndrome in rheumatoid arthritis. *Rheumatology* 2013;52:1084–9.
- 5 Seror R, Ravaud P, Bowman SJ, *et al*. EULAR Sjögren's syndrome disease activity index: development of a consensus systemic disease activity index for primary Sjögren's syndrome. *Ann Rheum Dis* 2010;69:1103–9.
- 6 Moutsopoulos HM, Webber BL, Vlagopoulos TP, *et al*. Differences in the clinical manifestations of sicca syndrome in the presence and absence of rheumatoid arthritis. *Am J Med* 1979;66:733–6.