

# Interferon signature in systemic autoimmune diseases: what does it mean?

Gaetane Nocturne, Xavier Mariette 

**To cite:** Nocturne G, Mariette X. Interferon signature in systemic autoimmune diseases: what does it mean?. *RMD Open* 2022;**8**:e002687. doi:10.1136/rmdopen-2022-002687

Received 7 October 2022  
Accepted 4 November 2022



© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

Department of Rheumatology, Université Paris-Saclay, INSERM UMR1184 : Center for immunology of viral infections and autoimmune diseases, Assistance Publique-Hôpitaux de Paris (AP-HP), Hôpital Bicêtre, Le Kremlin-Bicêtre, France

**Correspondence to**  
Dr Xavier Mariette;  
xavier.mariette@aphp.fr

The first description of the involvement of interferons (IFN) in systemic autoimmune diseases (AID) dates back more than 40 years.<sup>1</sup> IFNs play a crucial role in numbers of immunological pathways involve in AID, such as induction of dendritic cell (mDC), MHC expression, cytokines like BAFF, IL-2, IL-7. The aim of this editorial is not to discuss all these mechanisms linking IFNs to autoimmunity but to discuss the origin and signification of what is called ‘the IFN signature’ and how measuring it. The diseases in which this signature plays a prominent role are systemic erythematosus lupus (SLE), Sjögren’s syndrome (SS),<sup>2,3</sup> inflammatory myositis<sup>4</sup> and scleroderma.<sup>5</sup> The study of this IFN signature is still attracting the interest of research teams around the world, and for good reason: the advent of IFN-targeted therapies could revolutionise the outcome of patients with IFN-mediated diseases. However, behind this so-called IFN signature, there are still many grey areas: what type(s) of IFNs are we talking about? Who are the producers of IFNs? What is the origin of this IFN signature? What tools for assessing the IFN signature? Several techniques have been developed including IFN-inducible gene signatures; IFN-inducible proteins such as MxA, Galectin 9, Siglec1, IP-10; reporter system with WISH cells; dosage of the different subtypes of circulating IFNs. And finally, what are the consequences of this IFN signature in clinical practice, both in the classification of patients with the aim of achieving personalised medicine, and in the management of treatments, particularly at a time when therapies targeting IFNs are being developed?

The so-called signature corresponds to the evidence of an upregulation of transcripts induced by the different IFN subtypes: type I (IFN- $\alpha$  (of which there are 13 subtypes),  $\beta$ ,  $\epsilon$ ,  $\kappa$  and  $\omega$ ), type II (IFN- $\gamma$ ) and type III (IFN- $\lambda$  1

to 4). However, the limitation of this transcriptomic definition was because until recently it did not allow differentiation between the three families of IFNs and only IFN- $\gamma$  could be easily measured. The IFN signature is classically assessed by the level of expression of different mRNA induced by IFN. Several combinations of genes have been proposed to calculate IFN score. A five-gene signature (IFI44, IFI44L, IFIT1, IFIT3, MxA) has been proposed first.<sup>6</sup> A four-gene set (IFI27, IFI44, IFI44L and RSAD2) was applied for the analysis of the anti-IFNAR therapeutic antibody—anifrolumab—trials.<sup>7</sup> This score allows just a differentiation into IFN score high and low. Moreover, the transcriptomic overlap between the three types of IFNs, notably between type I and type III, is still a limit.<sup>8</sup> It is the reason why the development of new techniques for the determination of type I IFN such as the single molecule array (Simoa),<sup>9</sup> mesoscale diffusion technique and new methods of analysing the transcriptomic signature have changed the situation. Importantly, it became apparent that within a single disease, different IFN profiles could coexist. In SLE, it was shown that the IFN signature was not limited to IFN- $\alpha$ , but involved the gradual activation of three distinct modules underpinned by various IFNs, including type II.<sup>10</sup> In SS, a multiomic study was able to define four clusters of patients.<sup>11</sup> Among them, three seem to be driven by IFNs: cluster C1 corresponds to the strongest IFN type I and type II signature, cluster C3 to an intermediate IFN signature associated with B-cell activation and finally cluster C4 would be mainly linked to IFN type II. More recently, it has been shown that during SS, in the blood, it is IFN- $\alpha$  that drives this IFN signature. The level of circulating IFN- $\alpha$  was also associated with the clinical and biological characteristics of SS patients and with the prognosis, with

an association between the baseline IFN- $\alpha$  level and the development of systemic complications within 5 years.<sup>12</sup> Lastly, comparing the serum dosage of type I and type II IFN with the IFN signature will allow knowing if the high IFN signature is due to a high level of IFN, to an increased activation of pattern recognition receptors, or to an excessive response of IFN-responsive genes to IFN.

The cellular origin of the different IFN subtypes has also been re-evaluated. Initially, the focus was on plasmacytoid dendritic cells (pDCs), which are 'professionals' IFN- $\alpha$  secretors. However, the role of resident cells, keratinocytes, renal tubular cells, salivary gland epithelial cells appears to be important both in the early and late phases of SLE and SS.<sup>13 14</sup> In SLE, it has been shown that the IFN signature in tubular cells is associated with proliferative histology and markers of fibrosis, and IFN signature in keratinocytes is associated with clinical non-response to therapy.<sup>15</sup> In addition, in SLE, the role of neutrophils, via NETosis, has also been demonstrated in inducing IFN signature.<sup>16</sup>

Lastly, a key question is what the origin of this IFN signature is. Genetic, epigenetic and environmental factors may be involved. The role of genetic factors and in particular single nucleotide polymorphisms located on IFN pathways is well established.<sup>17 19</sup> In SLE, more than 100 genetic risk loci have been identified. Half of them are related to IFN pathways.<sup>20</sup> Interestingly, type I and type II are involved. SNP within interferon regulatory factor (IRF) 5 locus has been shown to be associated with SLE and SS.<sup>20 21</sup> Among these SNPs a 5-bp CGGGG indel in the promoter of *IRF5* is associated with a high level of *IRF5* mRNA, mainly after viral infection, which in turn correlate with higher IFN signature.<sup>22</sup> SNPs in *STAT4* loci are also associated with both SLE and SS.<sup>13 23</sup> Interestingly, *STAT4* is a classical transcription factor of type 2 IFN but an association between *STAT4* mRNA level and PKR, *IFITM1* and *MX1* mRNA levels that are type 1 IFN-induced genes has been described.<sup>24</sup> Epigenetic factors also play a role. Interestingly, epigenetic control seems to override genetic susceptibility. This has been demonstrated with methylation notably in SS where methylation alterations in B cells are more frequent in some specific pathways including IRF genes.<sup>25</sup> The most common pathological situation where type I IFN is increased is viral infections. A viral origin of systemic AID has been looked for decades. But even if some associations with a higher immune response to some viruses (especially Epstein-Barr virus<sup>26</sup>) have been observed in SLE and SS, no causal virus was found. If an exogenous viral infection cannot be found as the cause of this IFN signature, it might be useful to consider the possible role of endogenous viruses. Human endogenous retroviruses (HERVs) represent almost 10% of our genome.<sup>27</sup> They are exposed to epigenetic regulation and have been shown to be differentially expressed in patients with AID including multiple sclerosis or SLE. Differentially expressed HERVs are in close proximity

with IFN-induced genes and could modulate their expression.<sup>28</sup>

Besides these viral endogenous sequences, a number of long non-coding RNAs (lncRNAs) are present in our genome, more or less expressed and having a regulatory role by specific interactions with DNA, RNA and proteins.<sup>29</sup> In this issue of RMD, Joachims *et al* focus on the role of long lncRNAs as regulators of the IFN response in SS patients. They assessed blood transcriptome of SS patients looking at RNA-seq and comparing patients with and without anti-SSA/Ro60 auto-antibodies. They confirmed that the IFN signature is a characteristic feature of anti-SSA/Ro60+ patients. They assessed transcripts associated with IFN signature and found that 16 lncRNAs were strongly correlated with the IFN signature. Interestingly, through functional experiments, they showed that four antisense lncRNAs (NRIR, OAS123-AS1, MX1-AS1 and GBP5-AS1) were rapidly induced in response to IFN and that their increased expression acted as a massive upregulation signal to induce the expression of mRNAs. This study therefore demonstrates the potential role of these lncRNAs as targets and regulators of the IFN response, which may therefore participate in the amplification of this response during systemic AID.

Thus, our understanding of the origin and the consequences of the IFN signature is progressing. The question arises as to what this means in clinical practice. The development of therapies that target IFN (JAK inhibitors, anifrolumab) is an important step for the management of patients suffering from systemic AID. It was during the studies evaluating anifrolumab in SLE that the use of IFN signature monitoring in patients was developed. By pooling data from the TULIP trials into a post hoc study, it was shown that anifrolumab was more effective in the subgroup of SLE patients with high IFN signature.<sup>30</sup> Of note, the presence of serological biomarkers (anti-DNA, low C3, low C4) also allowed to discriminate patients who could benefit from anifrolumab treatment. In practice, it will therefore be interesting to compare the interest of monitoring the IFN signature (transcriptomic signature including 4 or 5 genes or inducible proteins) or the dosage of serum IFN itself by sensitive techniques (SIMOA or meso scale diffusion) versus the use of classical biomarkers more easily available in clinical practice. In addition, it will be important to assess the reproducibility of calculation of the IFN score. Actually, the IFN score calculated at different laboratories may be hardly comparable due to the distinct sets of IFN-stimulated genes assessed and different controls. The value of the IFN signature for stratification of patients treated with tsDMARDs is also being studied. JAK inhibitors may inhibit both type I by inhibiting JAK1 and TYK2 downstream IFNAR and also Type II IFN by inhibiting JAK1 and JAK2 downstream IFN gamma receptor. It was confirmed that they may decrease the IFN signature.<sup>31</sup> However, the IFN

signature at baseline has never been shown to predict clinical response to JAK inhibitors until recently. In a study presented at the 2022 ACR meeting focused on SS patients, it was shown that IFN signature at baseline correlated with response to filgotinib (JAK inhibitor) and to tirabrutinib (BTK inhibitor).<sup>32</sup> Finally, the evaluation of the IFN signature as a predictor of response to treatment could be also of interest in AID in which IFN is not the main driver. This is the case in rheumatoid arthritis (RA) in which an IFN signature is observed in only 20% of patients.<sup>33</sup> A recent study showed that in patients with early RA, an IFN signature, linked to IFN- $\alpha$ , was associated with a lower probability of response to csDMARD treatment at 6 months.<sup>33</sup> This was independent of conventional markers of disease activity. It was suggested that IFNs could drive disease persistence epigenetic reprogramming of autoimmune lymphocytes.

In conclusion, the understanding of the IFN signature in systemic AID and its usefulness in clinical practice continues to progress. Nevertheless, there are still challenges to overcome for understanding the origin of this IFN signature in both blood and tissue and the role of non-coding RNA sequences, deciphering the specificity of this signature regarding type I, type II or type III IFN. It will be needed to compare the IFN signature to the dosages of IFN themselves to determine what is the most relevant and the most convenient in daily practice and to explore the usefulness of the IFN signature compared with the usual biomarkers, notably autoantibodies for stratifying patients and optimising therapeutic strategies. Clarifying these points will help optimise the management of patients with numbers of systemic AIDs in which IFN play a role.

**Contributors** GN and XM wrote the editorial.

**Funding** The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

**Competing interests** None declared.

**Patient consent for publication** Not applicable.

**Ethics approval** Not applicable.

**Provenance and peer review** Commissioned; externally peer reviewed.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

**ORCID iD**

Xavier Mariette <http://orcid.org/0000-0002-4244-5417>

## REFERENCES

- 1 Hooks JJ, Moutsopoulos HM, Geis SA, *et al*. Immune interferon in the circulation of patients with autoimmune disease. *N Engl J Med* 1979;301:5–8.
- 2 Hall JC, Baer AN, Shah AA, *et al*. Molecular Subsetting of Interferon Pathways in Sjögren's Syndrome. *Arthritis Rheumatol* 2015;67:2437–46.
- 3 Gottenberg J-E, Cagnard N, Lucchesi C, *et al*. Activation of IFN pathways and plasmacytoid dendritic cell recruitment in target organs of primary Sjögren's syndrome. *Proc Natl Acad Sci U S A* 2006;103:2770–5.
- 4 Walsh RJ, Kong SW, Yao Y, *et al*. Type I interferon-inducible gene expression in blood is present and reflects disease activity in dermatomyositis and polymyositis. *Arthritis Rheum* 2007;56:3784–92.
- 5 Brkic Z, van Bon L, Cossu M, *et al*. The interferon type I signature is present in systemic sclerosis before overt fibrosis and might contribute to its pathogenesis through high BAFF gene expression and high collagen synthesis. *Ann Rheum Dis* 2016;75:1567–73.
- 6 Bodewes ILA, Al-Ali S, van Helden-Meeuwse CG, *et al*. Systemic interferon type I and type II signatures in primary Sjögren's syndrome reveal differences in biological disease activity. *Rheumatology* 2018;57:921–30.
- 7 Furie R, Khamashta M, Merrill JT, *et al*. Anifrolumab, an Anti-Interferon- $\alpha$  receptor monoclonal antibody, in moderate-to-severe systemic lupus erythematosus. *Arthritis Rheumatol* 2017;69:376–86.
- 8 Lazear HM, Schoggins JW, Diamond MS. Shared and distinct functions of type I and type III interferons. *Immunity* 2019;50:907–23.
- 9 Rodero MP, Decalf J, Bondet V. Detection of interferon alpha protein reveals differential levels and cellular sources in disease. Ultrasensitive detection of IFN $\alpha$  in human diseases. *J Exp Med* 2017;214:1547–55.
- 10 Chiche L, Jourde-Chiche N, Whalen E, *et al*. Modular transcriptional repertoire analyses of adults with systemic lupus erythematosus reveal distinct type I and type II interferon signatures. *Arthritis Rheumatol* 2014;66:1583–95.
- 11 Soret P, Le Dantec C, Desvaux E, *et al*. A new molecular classification to drive precision treatment strategies in primary Sjögren's syndrome. *Nat Commun* 2021;12:3523.
- 12 Trutschel D, Bost P, Mariette X. Variability in primary Sjögren's syndrome is driven by interferon alpha, and genetically associated with the class II HLA DQ locus. *Arthritis Rheumatol* 2022.
- 13 Psarras A, Wittmann M, Vital EM. Emerging concepts of type I interferons in SLE pathogenesis and therapy. *Nat Rev Rheumatol* 2022;18:575–90.
- 14 Rivière E, Pascaud J, Virone A. Interleukin-7/Interferon axis drives T-cell and salivary gland epithelial cell interactions in Sjögren's syndrome. *Arthritis Rheumatol* 2020.
- 15 Der E, Ranabothu S, Suryawanshi H, *et al*. Single cell RNA sequencing to dissect the molecular heterogeneity in lupus nephritis. *JCI Insight* 2017;2:93009.
- 16 Garcia-Romo GS, Caielli S, Vega B, *et al*. Netting neutrophils are major inducers of type I IFN production in pediatric systemic lupus erythematosus. *Sci Transl Med* 2011;3:73ra20.
- 17 Khatri B, Tessner KL, Rasmussen A, *et al*. Genome-Wide association study identifies Sjögren's risk loci with functional implications in immune and glandular cells. *Nat Commun* 2022;13:4287.
- 18 International Consortium for Systemic Lupus Erythematosus Genetics (SLEGEN), Harley JB, Alarcón-Riquelme ME, *et al*. Genome-wide association scan in women with systemic lupus erythematosus identifies susceptibility variants in ITGAM, PXX, KIAA1542 and other loci. *Nat Genet* 2008;40:204–10.
- 19 Dieudé P, Wipff J, Guedj M, *et al*. BANK1 is a genetic risk factor for diffuse cutaneous systemic sclerosis and has additive effects with IRF5 and STAT4. *Arthritis Rheum* 2009;60:3447–54.
- 20 Wang Y-F, Zhang Y, Lin Z, *et al*. Identification of 38 novel loci for systemic lupus erythematosus and genetic heterogeneity between ancestral groups. *Nat Commun* 2021;12:772.
- 21 Lessard CJ, Li H, Adrianto I, *et al*. Variants at multiple loci implicated in both innate and adaptive immune responses are associated with Sjögren's syndrome. *Nat Genet* 2013;45:1284–92.
- 22 Miceli-Richard C, Gestermann N, Itah M, *et al*. The CGGGG insertion/deletion polymorphism of the IRF5 promoter is a strong risk factor for primary Sjögren's syndrome. *Arthritis Rheum* 2009;60:1991–7.
- 23 Nocturne G, Mariette X. Advances in understanding the pathogenesis of primary Sjögren's syndrome. *Nat Rev Rheumatol* 2013;9:544–56.
- 24 Gestermann N, Mekinian A, Comets E, *et al*. Stat4 is a confirmed genetic risk factor for Sjögren's syndrome and could be involved in type 1 interferon pathway signaling. *Genes Immun* 2010;11:432–8.
- 25 Miceli-Richard C, Wang-Renault S-F, Boudaoud S, *et al*. Overlap between differentially methylated DNA regions in blood B lymphocytes and genetic at-risk loci in primary Sjögren's syndrome. *Ann Rheum Dis* 2016;75:933–40.
- 26 Harley JB, Harley ITW, Guthridge JM, *et al*. The curiously suspicious: a role for Epstein-Barr virus in lupus. *Lupus* 2006;15:768–77.

- 27 Tokuyama M, Kong Y, Song E, *et al.* ERVmap analysis reveals genome-wide transcription of human endogenous retroviruses. *Proc Natl Acad Sci U S A* 2018;115:12565–72.
- 28 Ng KW, Attig J, Bolland W, *et al.* Tissue-Specific and interferon-inducible expression of nonfunctional ACE2 through endogenous retroelement co-option. *Nat Genet* 2020;52:1294–302.
- 29 Statello L, Guo C-J, Chen L-L, *et al.* Gene regulation by long non-coding RNAs and its biological functions. *Nat Rev Mol Cell Biol* 2021;22:96–118.
- 30 Vital EM, Merrill JT, Morand EF, *et al.* Anifrolumab efficacy and safety by type I interferon gene signature and clinical subgroups in patients with SLE: post hoc analysis of pooled data from two phase III trials. *Ann Rheum Dis* 2022;81:951–61.
- 31 Dörner T, Tanaka Y, Petri M. 185 Baricitinib-associated changes in type I interferon gene signature during a 24-week phase 2 clinical SLE trial. *Lupus Sci Med* 2019;6.
- 32 Stratification on Baseline Interferon Scores and Use of CRESS to Improve the Design of Future Trials in Sjögren's Syndrome: Post Hoc Analysis from a Randomized Controlled Trial. ACR Meeting Abstracts. Available: <https://acrabstracts.org/abstract/stratification-on-baseline-interferon-scores-and-use-of-cress-to-improve-the-design-of-future-trials-in-sjogrens-syndrome-post-hoc-analysis-from-a-randomized-controlled-trial/> [Accessed 4 Oct 2022].
- 33 Cooles FAH, Tarn J, Lendrem DW, *et al.* Interferon- $\alpha$ -mediated therapeutic resistance in early rheumatoid arthritis implicates epigenetic reprogramming. *Ann Rheum Dis* 2022;10.1136/annrheumdis-2022-222370. [Epub ahead of print: 09 Jun 2022].