

Supplementary table 2: Characteristics of patients with radiographic score separated by diagnosis and those who fulfill modified New York criteria (radiographic) and patients who do not fulfill these criteria (non-radiographic)

	axSpA		axPsA	
	radiographic	non-radiographic	radiographic	non-radiographic
n	67	29	37	142
Female gender, n (%)	21 (31)	16 (55)	14 (38)	77 (54)
Age, mean (SD)	46.9 (13.7)	47.4 (12.6)	56.5 (8.4)	50.5 (12.1)
Symptom duration, mean (SD)	18.9 (12.5)	12 (10)	16.5 (12.6)	8.3 (6.6)
Disease duration, mean (SD)	11.6 (11.3)	4.9 (6.3)	10.7 (9.9)	5.6 (5.6)
HLA B-27 positive, n (%)	55 (85)	17 (65)	13 (42)	19 (17)
CRP mg/l, mean (SD)	13 (19)	12.1 (19.3)	7 (6.4)	7.6 (11.3)
CRP >=5 mg/L, n (%)	31 (49)	16 (57)	18 (51)	55 (41)
BMI, mean (SD)	27.4 (5.7)	28.6 (6.4)	30 (6.4)	29 (6.4)
Obesity (BMI>=30), n (%)	21 (32)	8 (30)	14 (40)	53 (38)
peripheral manifestations, n (%)	14 (21)	16 (55)	32 (86)	125 (88)
Uveitis, ever, n (%)	18 (27)	2 (7)	2 (5)	4 (3)
IBD, ever, n (%)	5 (7)	3 (10)	4 (11)	1 (1)
physician global disease activity (NRS), mean (SD)	5.9 (2.4)	5.5 (2.1)	5.1 (2)	5.1 (1.9)
patient global disease activity (NRS), mean (SD)	5.5 (2.5)	6.1 (2.4)	5.8 (2.1)	5.6 (2.3)
patient pain (NRS), mean (SD)	5.5 (2.5)	6.1 (2.4)	5.9 (2.1)	5.3 (2.4)
Sleep disorder (NRS), mean (SD)	5.7 (3)	6.1 (3.2)	6.1 (2.8)	5.2 (3.1)