

SUPPLEMENTARY DATA

Supplementary Table S2. Serologic data for the patients with early rheumatoid arthritis

	Patients (n = 566)	Patients without RRP at 1 year (n = 421)	Patients with RRP at 1 year (n = 145)	p value
Serologic data				
Anti-CCP2 (cut-off: 50 U/mL), no. (%)				0.0001
negative	301 (53.2)	246 (58.4)	55 (37.9)	
low	44 (7.8)	29 (6.9)	15 (10.3)	
high	221 (39.0)	146 (34.7)	75 (51.7)	
Anti-CCP2 (cut-off: 40 U/mL; Sp 98%), no. (%)				< 0.0001
negative	300 (53.0)	246 (58.4)	54 (37.2)	
low	28 (4.9)	17 (4.0)	11 (7.6)	
high	238 (42.0)	158 (27.5)	80 (55.2))	
Anti-MCV (cut-of: 35 UA/mL; Sp 98%), no. (%)				< 0.0001
negative	297 (52.5)	242 (57.5)	55 (37.9)	
low	42 (7.4)	32 (7.6)	10 (6.9)	
high	227 (40.1)	147 (34.9)	80 (55.2)	
AhFibA (cut-off 0.119 AU; Sp 98%), no. (%)				< 0.0001
negative	291 (51.4)	237 (56.3)	54 (37.2)	
low	43 (7.6)	28 (6.6)	15 (10.3)	
high	232 (41.0)	156 (37.1)	76 (52.4)	

Sp: diagnostic specificity for rheumatoid arthritis