

Supplementary table 1: Propensity-score for the first and second TNFi derived weights per visit

	<i>Min</i>	<i>Median [IQR]</i>	<i>Mean (S.D.)</i>	<i>Max</i>	<i>N patients at risk.</i>
First TNFi					
<i>Month 1</i>	1.01	1.05[1.03-1.10]	2.01(4.76)	53.44	708
<i>Month 6</i>	1.06	1.13[1.10-1.22]	2.00(2.46)	14.50	627
<i>Month 12</i>	1.02	1.06[1.04-1.10]	1.99(4.11)	39.18	520
<i>Month 18</i>	1.00	1.03[1.01-1.05]	2.00(6.24)	73.26	469
<i>Month 24</i>	1.01	1.04[1.02-1.04]	2.01(5.57)	51.43	430
<i>Month 36</i>	1.01	1.04[1.02-1.09]	2.02(5.08)	57.06	378
<i>Month 48</i>	1.02	1.05[1.04-1.09]	1.99(4.29)	33.69	322
<i>Month 60</i>	1.01	1.03[1.02-1.06]	1.99(5.47)	52.19	272
Second TNFi					
<i>Month 6</i>	1.01	1.05[1.02-1.09]	2.03(5.54)	71.41	258
<i>Month 12</i>	1.02	1.09[1.05-1.17]	1.98(3.64)	40.92	241
<i>Month 18</i>	1.03	1.07[1.01-1.16]	1.99(3.34)	21.87	216
<i>Month 24</i>	1.00	1.10[1.08-1.14]	2.00(3.19)	25.92	193
<i>Month 36</i>	1.00	1.10[1.04-1.25]	2.05(4.05)	33.14	171
<i>Month 48</i>	1.03	1.12[1.07-1.20]	1.12(2.88)	20.57	144
<i>Month 60</i>	1.01	1.94[1.03-1.20]	1.94(4.00)	37.22	121

Footnote: IQR= interquartile range. SD= standardized deviation. TNFi= Tumor Necrosis Factor inhibitor.