T1w/CT	CT+	CT-	Total	SE	0.90	0.60 to 0.99
T1w+	9	7	16	SP	0.73	0.54 to 0.86
T1w-	1	19	20	DA	0.78	0.61 to 0.90
Total	10	26	36	PPV	0.56	0.33 to 0.77
				NPV	0.95	0.76 to 1.00
VIBE/CT	CT+	CT-	Total	SE	1.00	0.72 to 1.00
VIBE+	10	3	13	SP	0.88	0.71 to 0.96
VIBE-	0	23	23	DA	0.92	0.77 to 0.98
Total	10	26	36	PPV	0.77	0.50 to 0.92
				NPV	1.00	0.86 to 1.00
SWI/CT	CT+	CT-	Total	SE	0.80	0.49 to 0.96
SWI+	8	1	9	SP	0.96	0.81 to 1.00
SWI-	2	25	27	DA	0.92	0.78 to 0.98
Total	10	26	36	PPV	0.89	0.57 to 0.99
				NPV	0.93	0.77 to 0.99

Supplementary Table 2: Results of contingency table analysis on per-metacarpophalangeal level. SWI = susceptibility-weighted imaging, VIBE = volumetric interpolated breath-hold examination, T1w = T1-weighted imaging sequence, CT = computed tomography, SE = sensitivity, SP = specificity, DA = diagnostic accuracy, PPV = positive predictive value, NPV = negative predictive value.