## Supplemental File - COVID-19 outcomes and treatment

NIH COVID-19 ordinal scale <sup>1</sup>	Mild <sup>2</sup>	Moderate	Severe <sup>3</sup>
	Score 1-3	Score 4-5	Score 6-8
All patients	11	0	1
Patients who received COVID-19 treatment <sup>1</sup>	9	0	0
Monocionals*	8	0	0
Oral antivirals <sup>+</sup>	2	0	0

<sup>&</sup>lt;sup>1</sup>National Institutes of Health (NIH) COVID-19 ordinal scale: 1) Not hospitalized and no limitations of activities

- 2) Not hospitalized, with limitation of activities, home oxygen (O2) requirement, or both
- 3) Hospitalized, not requiring supplemental O2 and no longer requiring ongoing medical care
- 4) Hospitalized, not requiring supplemental O<sub>2</sub> but requiring ongoing medical care
- 5) Hospitalized, requiring any supplemental O<sub>2</sub>
- 6) Hospitalized, requiring noninvasive ventilation or use of high-flow O<sub>2</sub> devices
- 7) Hospitalized, receiving invasive mechanical ventilation or extracorporeal membrane oxygenation
- 8) Death

- \*7 Bebtelovimab, 1 sotrovimab
- + Nirmatrelvir/ritonavir

<sup>&</sup>lt;sup>2</sup>Score 1 = 5 patients, Score 2 = 6 patients

<sup>&</sup>lt;sup>3</sup>Score 6 = 1 patient

<sup>¶1</sup> patient received both a monoclonal infusion and oral treatment